Astronomy & Astrophysics manuscript no. 09595
April 14, 2008
© ESO 2008

arXiv:0804.2184v1 [astro-ph] 14 Apr 2008

Young stars and brown dwarfs surrounding Alnilam (ϵ Ori) and Mintaka (δ Ori)

J. A. Caballero[1,2] and E. Solano[3]

[1] Dpto. de Astrofísica y Ciencias de la Atmósfera, Facultad de Física, Universidad Complutense de Madrid, E-28040 Madrid, Spain
[2] Max-Planck-Institut für Astronomie, Königstuhl 17, D-69117 Heidelberg, Germany
[3] Laboratorio de Astrofísica Espacial y Física Fundamental, INSA, Apdo. 78, E-28691 Villanueva de la Cañada, Madrid, Spain

Received 17 February 2008; accepted 9 April 2008

ABSTRACT

Aims. We look for new regions for the search of substellar objects.
Methods. Two circular areas, 45 arcmin-radius each, centred on the young massive star systems Alnilam and Mintaka in the Orion Belt, have been explored. The regions are very young (less than 10 Ma), have low extinction, and are neighbours to σ Orionis (~3 Ma), a young open cluster very rich in brown dwarfs and planetary-mass objects. We have used Virtual Observatory tools, the astro-photometric Tycho-2, DENIS and 2MASS catalogues, 10 control fields at similar galactic latitudes, and X-ray, mid-infrared and spectroscopic data from the literature.
Results. We have compiled exhaustive lists of known young stars and new candidate members in the Ori OB1b association, and of fore- and background sources. A total of 136 stars display features of extreme youth, like early spectral types, lithium in absorption, or mid-infrared flux excess. Other two young brown dwarf and 289 star candidates have been identified from an optical/near-infrared colour-magnitude diagram. We list additional 74 known objects that might belong to the association. This catalogue can serve as an input for characterisation of the stellar and high-mass substellar populations in the Orion Belt. Finally, we have investigated the surface densities and radial distributions of young objects surrounding Alnilam and Mintaka, and compared them with those in the σ Orionis cluster. We report a new open cluster centred on Mintaka.
Conclusions. Both regions can be analogs to the σ Orionis cluster, but more massive, more extended, slightly older, and less radially concentrated.

Key words. astronomical data bases: miscellaneous – stars: low mass, brown dwarfs – open clusters and associations: individual: Ori OB1b, Collinder 70 – stars: individual: Alnilam, Mintaka

1. Introduction

The knowledge of the frequency and characteristics of brown dwarfs (substellar objects with masses below the hydrogen burning limit) is essential for the most advanced scenarios of fragmentation of molecular clouds and very low-mass star formation (Reipurth & Clarke 2001; Bate, Bonnell & Bromm 2003; Whitworth et al. 2007). In particular, they provide valuable information on the bottom of the Initial Mass Function (e.g. Luhman et al. 2000). Brown dwarfs are found in the field, both as companions to stars (Nakajima et al. 1995; Rebolo et al. 1998; Goldman et al. 1999) and free-floating (Delfosse et al. 1997; Ruiz, Leggett & Allard 1997; Kirkpatrick et al. 1999). Brown dwarfs are much brighter when younger (Chabrier & Baraffe 2000); they are, thus, very common in young open clusters and star-forming regions, such as the Pleiades (Rebolo, Zapatero Osorio & Martín 1995), ρ Ophiuchi (Luhman, Liebert & Rieke 1997), Chamaeleon I+II (Neuhäuser & Comerón 1998), Taurus-Auriga (Briceño et al. 1998) or the Orion Nebula Cluster (Hillenbrand & Carpenter 2000). There are, however, limitations in the search for brown dwarfs in these regions and in others: in the youngest ones (e.g. Chamaeleon, Ophiuchus), there is variable extinction that hinders the characterisation of the recently-born brown dwarfs, while in the others (e.g. Pleiades, Hyades), the relatively old brown dwarfs have dimmed down to faint magnitudes that force to use very large, expensive, astronomical facilities.

There exists, nevertheless, a cornerstone for the search of brown dwarfs and objects below the deuterium burning-mass limit: the σ Orionis cluster in the Ori OB1b association (Garrison 1967; Lyngå 1981; Walter et al. 1997). The cluster is very young (~3 Ma), practically free of extinction ($A_V \lesssim 0.3$ mag) and relatively nearby ($d \sim 385$ pc). See Caballero (2007a, 2008c) for extensive bibliographic and data compilations on the cluster. It does not only harbour a rich population of OB-type, Herbig Ae/Be and T Tauri stars, but also Herbig-Haro objects, X-ray emitters and substellar objects (Wolk 1996; Reipurth et al. 1998; Oliveira & van Loon 2004; Franciosini,

Send offprint requests to: José Antonio Caballero, e-mail: caballero@astrax.fis.ucm.es

Pallavicini & Sanz-Forcada 2006). Indeed, the σ Orionis cluster possesses the best spectroscopically investigated and most numerous population of brown dwarfs and planetary-mass objects down to a few Jupiter masses (Zapatero Osorio et al. 2000; Béjar et al. 2001; Caballero et al. 2006). Besides, an important fraction of the σ Orionis area has been already covered or is being currently investigated by very deep, wide photometric surveys, screening the whole brown dwarf and part of the planetary mass regimes (González-García et al. 2006; Caballero et al. 2007; Bihain et al., in prep.).

To compare substellar mass functions, spatial distributions or disc frequencies of different clusters, and to look for new brown dwarfs and planetary-mass objects, it is necessary, therefore, to search for new locations. Youth, closeness and low extinction, just like in σ Orionis, are strongly required. Since the new hunting grounds for the search of substellar objects must resemble σ Orionis, it is natural to look for them not far away, just in the Ori OB1b association.

1.1. Alnilam and Mintaka

A "clustering of early-type stars elongated roughly parallel to the Galactic plane" (Guetter 1979) was firstly noticed by Pannekoek (1929). Later, in his classical review of nearby O-type associations, Blaauw (1964) described an Ori OB1 complex splitted into four divisions, being Ori OB1b (the "Orion Belt") one of them. Wide survey observations with Schmidt telescopes have shown a patent overdensity of Hα emission stars in the area (Haro & Moreno 1953; Wiramihardja et al. 1989). Canonical age and heliocentric distance are in the intervals 1–7 Ma and 350–500 pc (e.g. Anthony-Twarog 1982; Lyngå 1987; Blaauw 1991; Brown et al. 1994; de Zeeuw et al. 1999; Harvin et al. 2002). The most representative stars in the Ori OB1b association are the bright O-type supergiants Alnitak, Alnilam and Mintaka (ζ Ori, ϵ Ori and δ Ori, respectively), that constitute the celebrated asterism of the Orion Belt. At least one star of the trio (Alnilam) was depicted in the Farnese Atlas and, therefore, tabulated in the original Hipparchus catalogue (Schaefer 2005). The easternmost supergiant, Alnitak, is nearly embedded in the core of the L1630/Orion B molecular cloud complex, which contains, among other nebulosities and H II regions, the Flame Nebula (NGC 2024) and the Horsehead Nebula. The high variable extinction and emission in the area (Jaffe et al. 1994; Lacy et al. 1994; Kramer, Stutzki & Winnewisser 1996) prevent from suitably studying its stellar and substellar populations *à la* σ Orionis.

The stellar populations in the Ori OB1b association have been characterised after Blaauw's (1964) seminal work by many authors (Hardie, Hesser & Tolbert 1964; Warren & Hesser 1978; Guetter 1981; Brown et al. 1994; Hernández et al. 2005). In contrast to the hypothesis of Sharpless (1962), who claimed a lack of noticeable excess of members in Ori OB1b later than A5, at least two nearby clusters are known within the association: σ Orionis, centred on the eponym σ Ori Trapezium-like star system (see above), and Collinder 70, centred on Alnilam (Collinder 1931). There is an additional open cluster in the background, to the north of Mintaka: Berkeley 20. It is a unusual, low-metallicity, old open cluster at $d \sim 8.4$ kpc, and about 2.5 kpc below the Galactic plane (Lyngå 1987; MacMinn et al. 1994).

While the size of σ Orionis is well determined at 20–30 arcmin (Béjar et al. 2004; Sherry, Walter & Wolk 2004; Caballero 2008a), the actual size of Collinder 70 is not ascertained. Some authors (e.g. Gieseking 1983; Dias, Lépine & Alessi 2001) have identified the Collinder 70 cluster, sometimes called the "ϵ Orionis cluster", as the whole Ori OB1b association. Markarjan (1951) (and, therefore, Lyngå 1987) tabulated an angular diameter of 149 arcmin, which would make the cluster to comprise the stellar populations surrounding Mintaka and σ Ori. Subramaniam et al. (1995) proposed that both Collinder 70 and NGC 1981 (to the north of the Orion Nebula Cluster) form a "probable binary open star cluster in the Galaxy". Some classical works also catalogued a bright diffuse galactic nebulae centred on Alnilam, and with an apparent size of 50 arcmin (Dreyer 1888 – NGC 1990; Cederblad 1946 – Ced 55h). The existence of (numerous) small cometary globules, remnant molecular clouds and giant outflows close to Alnilam and Mintaka, even in larger amount than close to σ Ori (Cernicharo et al. 1992; Yun et al. 1997; Ogura & Sugitani 1998; Mader et al. 1999), favours the hypothesis of a wide region with a rather homogeneous age of no more than ~7 Ma. See also Wolk (1996) and Scholz & Eislöffel (2005) for other age determinations. The recent determinations of heliocentric distances to VV Ori AB, a double-lined eclipsing binary in a detached configuration close to Alnilam (d = 388±30 pc; Terrell, Munari & Siviero 2007), and to σ Ori AB, one of the most massive binaries known ($d \sim 385$ pc assuming the hierarchical triple scenario – Caballero 2008b; D. M. Peterson et al., in prep.), seem to support the homogeneity of the region. There are hints, nonetheless, of subestructure and overlapping populations within the association, as firstly noticed by Hardie et al. (1964) and Warren & Hesser (1977), and later by Jeffries et al. (2006), Caballero (2007a) and Caballero & Dinis (2008).

Searches for low-mass young stars in the central and western regions of the Ori OB1b association (the σ Orionis cluster is to the east) have been already carried out by Sherry, Walter & Wolk (2000), Sherry (2003) and Briceño et al. (2005). Several very low-mass star and brown dwarf candidates have also been identified surrounding Alnilam by Béjar (2001), Pérez-Garrido, Díaz-Sánchez & Villo (2005) and Scholz & Eislöffel (2005). These works are, however, biased towards very low-mass objects or incomplete. On the one hand, Pérez-Garrido et al. (2005) selected brown dwarf candidates in the 2MASS catalogue (Skrutskie et al. 2006) without optical counterpart in the USNO-A2.0 catalogue (Monet et al. 1998); the majority, if not all, of their 23 objects are, nonetheless, identified in the red optical passbands of the most recent, deeper USNO-B1.0 and DENIS catalogues (Epchtein et al 1999; Monet et al. 2003). On the other hand, Béjar (2001) and Scholz & Eislöffel (2005) selected their cluster member candidates from deep optical surveys (*IZ*, *RI*) in ~1000 arcmin^2-wide areas to the southeast and the northwest of Alnilam, respectively. Each of them might have surveyed less than one quarter of the minimum size of Collinder 70, and only for objects fainter than I = 14–16 mag.




Fig. 1. Pictogramme showing the Orion-Lepus region and the survey area. *Stars*: some of the brightest stars of Orion, Lepus and adjacent constellations (sizes are roughly proportional to brightness). The filled stars are Alnilam and Mintaka. *Circles*: comparison fields and the survey area around Alnilam and Mintaka. *Solid line*: parallel of constant declination $\delta = +2$ deg. There are no DENIS data northern (i.e. to the right) of the parallel. Colour versions of all our figures are available in the electronic publication.

Béjar et al. (2003a) and Béjar, Caballero & Rebolo (2003b) carried out the spectroscopic follow-up of nine "ϵ Orionis cluster" photometric member candidates presented in Béjar (2001). Derived spectral types ranged between M4.5 and M8, at the expected star-brown dwarf boundary. Some of the spectra also showed features of extreme youth (Li I λ6707.8 Å in absorption, Hα λ6652.8 Å in emission, and/or faint alkali lines – a signpost of low surface gravity, low g). No searches for brown dwarfs have been carried out in the Mintaka region yet.

In the present work, we compile lists of confirmed very young stars, association member candidates and fore- and background sources in two wide survey areas centred on Alnilam and Mintaka, using tools of the Virtual Observatory. The final catalogue, ranging from the two massive OB-type supergiants to intermediate M-type substellar objects, is useful for next studies on characterisation of stars and brown dwarfs, initial mass function, frequency and properties of discs and multiplicity in the Ori OB1b association, and complements past and future works (e.g. Béjar et al., in prep.; Pérez-Garrido et al., in prep.). Finally, we present a preliminar study of the spatial distribution of association members and candidates.

2. Analysis and results

2.1. Data and survey areas

The search for stellar and substellar members of the Ori OB1b association around Alnilam and Mintaka were conducted using the Tycho-2 (Høg et al. 2000), DENIS and 2MASS catalogues and photometric, spectroscopic and astrometric information from the literature. Two main survey fields were defined as

Table 1. Observed fields (45 arcmin-radius each).

Name	α (J2000)	δ (J2000)	l [deg]	b [deg]
Mintaka	05 32 00.40	–00 17 56.7	203.9	–17.7
Alnilam	05 36 12.81	–01 12 06.9	205.2	–17.2
c_1	05 59 46.51	–11 57 00.3	218.0	–16.8
c_2	05 55 49.69	–13 26 39.1	219.0	–18.3
c_3	06 03 04.36	–13 41 16.2	220.0	–16.8
c_4	05 59 03.14	–15 10 29.9	221.0	–18.3
c_5	06 06 20.78	–15 25 51.3	222.0	–16.8
c_6	06 02 15.03	–16 54 41.1	223.0	–18.3
c_7	06 09 36.11	–17 10 44.3	224.0	–16.8
c_8	06 05 25.67	–18 39 11.3	225.0	–18.3
c_9	06 12 50.69	–18 55 53.7	226.0	–16.8
c_{10}	06 08 35.37	–20 23 59.1	227.0	–18.3

45 arcmin-radius circles centred on the stars. Since the two circular fields are almost tangent, the size choice simultaneously allows exploring a large manageable area of about 3.5 deg^2, and prevents overlapping between them (there is, however, a tiny overlapping of a few arcmin^2 between both circular fields). The radius of investigation is roughly twice the radius of the core of the σ Orionis cluster, that is expected to be a model. To discard background and foreground objects in the two main fields, we selected ten nearby, same-size comparison fields, c_i (i = 1, 2... 10), fulfilling the following requirements:

- data availability (i.e. $\delta < +2$ deg; there is no DENIS data for larger declinations);
- separation from the Orion region to avoid gas clouds and dispersed young stellar populations;
- same galactic latitude to the main fields ($b \sim -17.5$ deg);
- similar galactic longitude to the main fields, but keeping away from the Galactic bulge to minimise the number of background giant stars ($l \sim$ 215–230 deg).

Central coordinates (equatorial and Galactic) of the twelve fields are given in Table 1. Both main and comparison fields were explored down to limiting magnitude V_T = 11.5 mag (Tycho-2), i = 18.0 mag (DENIS), J, H, K_s = 17.1, 16.4, 14.3 mag (2MASS) [in Tables A.3 and A.4, there are Tycho-2 stars fainter than the indicated V_T-band limiting magnitude]. A pictogramme of the survey areas is given in Fig, 1, while Fig. 2 shows the images of the fields surrounding Alnilam and Mintaka and of one comparison field.

In what follows, information retrieval, data manipulation, filtering and selection has been done taking advantage of Virtual Observatory[1] standards and tools, in particular Aladin[2] (Bonnarel et al. 2000) and TOPCAT[3]. The Virtual Observatory is an international, community-based initiative to provide seamless access to the data available from astronomical archive

[1] http://www.ivoa.net
[2] http://aladin.u-strasbg.fr/aladin.gml
[3] http://www.star.bris.ac.uk/ mbt/topcat/

and services, as well as to develop state-of-the-art tools for the efficient analysis of this huge amount of information. Padovani et al. (2004), Tsalmantza et al. (2006), and Caballero & Solano (2007) are examples of the efficiency of such tools in helping astronomers to produce scientific results.

2.2. Bright early-type stars

We have followed a procedure very similar to that carried out in Caballero (2007a) to identify bright very young stars using Tycho-2 and 2MASS astrometric and photometric data. First, we loaded the data within the twelve 45 arcmin-radius fields. Secondly, a cross-correlation between Tycho-2 and 2MASS was done using the Catalog Cross Match tool implemented in Aladin. For each Tycho-2 source, the 2MASS counterpart was defined as the nearest source found in a circle centred on the Tycho-2 source and radius of 4 arcsec (the default threshold). A total of 1276 Tycho-2 sources with 2MASS counterpart were identified within the ten comparison fields, which yielded $\overline{N}$ = 130 objects per field [$\sigma(N) = 30$]. The actual number of Tycho-2/2MASS sources per c_i field varies between 83 (c_2) and 169 (c_7), while there are $N = 111$ and 113 stars in the Alnilam and Mintaka fields, respectively. The B_T, V_T, J, H and K_s magnitudes of the 224 stars are provided in Tables A.1 and A.2, one for each main field.

Although the number N of Tycho-2/2MASS stars surrounding Alnilam and Mintaka does not *a priori* support the hypothesis of a stellar overdensity, there is an evident overdensity of bright *blue* stars surrounding the two supergiants (as expected from an OB association). In the ten comparison fields, there are only six stars bluer than $V_T - K_s = 0.0$ mag and brighter than $V_T = 10.0$ mag. Two of them have accurate determinations of the parallax ($\pi/\delta\pi > 3$) and are significantly closer to the Sun than the Ori OB1b association: HD 40071 (d = 250±70 pc) and HD 40355 AB (d = 180±30 pc). The other four stars (HD 41367 AB, HD 41488, HD 41737 and HD 42263) are B8–9 dwarfs, subdwarfs and subgiants located at tens of degrees from the young Orion associations that seem to populate the interstellar field. Therefore, ∼0.6 (between 0 and 1) interloper stars bluer than $V_T - K_s = 0.0$ mag and brighter than V_T = 10.0 mag are expected to be in each of the Alnilam and Mintaka fields. However, there are actually 17 and 11 such stars, respectively. Hence, there are between 20 and 30 more bright blue stars surrounding the two supergiants than in other regions at the same Galactic latitude and far from the Orion star-forming region. Given the Tycho-2 limiting magnitudes and the expected spectral types (i.e. colours) of young stars at $d \sim$ 385 pc, we could only detect an overdensity of blue (i.e. early-type) stars. This calculated overdensity is similar to that in the σ Orionis cluster (Caballero 2007a) and justifies next steps.

We have added other 21 Tycho-2/2MASS stars with blanks in the Tycho-2 proper motion that were not considered by the Aladin Catalog Cross Match tool to the list of 224 correlated Tycho-2/2MASS stars. In those cases, we have taken the proper motions from the USNO-B1 and Tycho-1 catalogues (Høg et al. 1998). This addition makes the sample of bright stars to increase up to 245.

With the help of optical-near infrared colour-magnitude diagrams, the Tycho-2 proper motions, *IRAS* fluxes, spectroscopic data from the literature and data from Vizier catalogues obtained using UCD[4]-based searches (Ochsenbein, Bauer & Marcout 2000), we have classified the stars in the Alnilam and Mintaka fields into three groups, separated by decreasing probability of membership in the Ori OB1b association. On the one hand, the classification is illustrated with the six panels in Fig. 3. On the other hand, Tables A.3 to A.8 show the results of the classification. The three groups are:

- stars with signposts of youth (Tables A.3 and A.4). By features of youth we understand early spectral types (O and B), Li I in absorption, *strong* X-ray or Hα emission (possibly associated to accretion processes), and infrared excesses by circumstellar material. We have also included low proper motion early A-type stars that follow the sequence defined by the remaining young stars in the colour-magnitude diagrams. There are only two stars with previously determined spectral type later than A: RX J0535.6–0152 AB and SS 28. Both of them display, however, features common of the T Tauri phase (see notes in Tables A.3 and A.4). We have also considered additional youth features, like the star being in the Herbig Ae/Be phase (HAeBe) or having a Vega-like disc.
- stars with unknown association membership status (Tables A.5 and A.6). They are stars with proper motions $\mu < 12$ mas a^{-1} that do not deviate very much from the young-star sequence in the colour-magnitude diagrams but have no known, clear, signposts of youth. Some A-type stars with $\mu \sim$ 5–12 mas a^{-1} that do not follow the sequence of the confirmed young stars are in this class. The threshold for the maximum μ of 12 mas a^{-1} is larger than in Caballero (2007a), who used $\mu > 10$ mas a^{-1} to identify stars with high tangential velocities among a list of photometric member candidates of the Ori OB1b association (the difference in the mean proper motion of the association, almost null, is not significant). The new conservative threshold allows us to recognize some bona-fide young stars with relatively large proper motions;
- stars that do not belong to the association (Tables A.7 and A.8). This class comprises: (*i*) foreground stars with proper motions $\mu > 12$ mas a^{-1}; (*ii*) foreground G-, K- and M-type stars with spectral type determination; (*iii*) *Hipparcos* stars with $\pi/\delta\pi > 3$ and heliocentric distances less than 250 pc; (*iv*) red stars without spectral type determination, and with colours $V_T - K_s > 2.5$ mag and spectral energy distributions of K–M-type stars; (*v*) very red objects with colours $V_T - K_s > 4.5$ mag and without flux excess due to discs in the mid-infrared (see notes in Appendix A). There are stars that simultaneously satisfy more than one criterion. For example, HD 36443 in the Mintaka field is a high proper motion ($\mu \approx 488$ mas a^{-1}) G5V star located at $d \sim 38$ pc and with a radial velocity $V_r \approx 9$ km s^{-1} discordant with association membership (Adams et al. 1929; Roman 1955; Perryman et al. 1997).

[4] Unified Content Descriptor.




Fig. 2. Inverted-colour, SERC B_J (photographic blue) DSS1 images of the Alnilam (left), Mintaka (centre) and c_1 (right) fields. Circles (in blue) indicate Tycho-2/2MASS stars. Approximate sizes are $1.5 \times 1.5\,\mathrm{deg}^2$. North is up, east is left. A tonge-shaped nebula appears in both Alnilam and Mintaka images (IC 434; see Fig. A.1).

Table 2. Number of bright stars per group and field.

Field	N members	N non-members	N unknown	N all
Alnilam	49	39	34	122
Mintaka	29	63	31	123
Total	78	102	65	245

Provided coordinates and proper motions in Tables A.3 to A.8 are from Tycho-2 (some exceptions are indicated). The majority of the spectral types and features of youth listed in the tables have been borrowed from the literature. The references are indicated in the last column. Abundant notes and remarks on the discussed stars are also provided in Appendix A. The most used works have been the spectroscopic studies in the Orion Belt by Guetter (1976, 1979) and the Henry Draper Extension Charts by Nesterov et al. (1995). The latter authors compiled positions, proper motions, photographic magnitudes and spectral types from the original works by Cannon and Cannon & Pickering (e.g. Cannon & Pickering 1918–1924). Table 2 summarizes the number of stars in each class and the total number of correlated Tycho-2/2MASS stars.

From the comparison with the c_i fields, between 0 and 1 late B-type stars surrounding Alnilam and Mintaka may actually be fore- or background B8–9-type stars. The contamination rate in the group of stars with signposts of youth is, therefore, ~3.5–5.5 % for this spectral types, and null for late O- and early B-type stars. On the other hand, from the number of stars in the comparison fields with colours $0.0\,\mathrm{mag} < V_T - K_\mathrm{s} < 0.5\,\mathrm{mag}$ (28 of the 1276 stars in the ten fields), it is expected that only ~3 early A-type stars in the foreground contaminate the Alnilam and Mintaka fields. It leads to estimate the contamination by such stars in Tables A.3 to A.6 at less than 10 %.

2.3. Intermediate- and late-type stars and brown dwarfs

We have performed a correlation between the DENIS and 2MASS catalogues identical to the Tycho-2/2MASS one. In this case, we have analysed the optical i (DENIS) and near-infrared JHK_s (2MASS) counterparts of more than 10^5 sources distributed amongst the ten comparison and the two Orion Belt fields. In particular, in the Alnilam and Mintaka fields, we have compiled the coordinates and four-band photometry of 10523 and 8288 sources, respectively.

A total of 50 DENIS/2MASS stars with spectroscopic features of youth (i.e. with Li I, Hα, low g; see Section 1.1) found in the literature have been identified in the Orion fields. Most of them are located surrounding Alnilam and come from wide prism-objective surveys of Hα emitters by, e.g., Haro & Moreno (1953; Haro objects) and Wiramihardja et al. (1989, 1991; Kiso objects), and from the spectroscopic analyses by Béjar et al. (2003a, 2003b; E Ori objects). We have not been able to identify V1299 Ori, an hypothetical B-type star close to the remnant molecular cloud [OS98] 40B. On the contrary, we have identified 8 DENIS/2MASS stars without known spectroscopic features of youth, but with *strong* X-ray emission detected by the *Einstein* and *ROSAT* satellites, that can be ascribed to a young age. Only three of them had been previously catalogued (MacDowell 1994; Ueda et al. 2001). Their colours are typical of young stars close to the main sequence. The results of *XMM-Newton* observations centred on Alnilam will be described in Caballero et al. (in prep.).

The names, coordinates, i- and K_s-band magnitudes, features of youth and references of the 58 young stars are provided in Tables A.9 and A.10. We list other spectro-photometric signposts of youth, such as Ca II $\lambda\lambda$3933.7,3968.5 Å (H and K lines), Hβ λ4861.3 Å, and [O I] λ6300.3 Å in emission (indicative of strong magnetic activity and/or outflows), and mid-infrared flux excess (mIR; suggestive of the presence of a circumstellar disc). Other 17 DENIS/2MASS stars in the literature have unreliable features of youth, such as faint Hα emission with $i - K_\mathrm{s}$ colour typical of field stars, conservative upper




Fig. 3. The Tycho-2/2MASS cross-match in the 10 comparison fields (top), and Alnilam (middle) and Mintaka (bottom) fields. Stars classified as association Ori OB1b non-members are marked with (blue) crosses. Remaining stars are marked with (blue) dots, except for stars with signposts of youth, that are marked with (red) filled stars. *Left panel*: proper motion diagrams. Big circles separate stars with proper motions larger than 12 mas a^{-1}. *Right panel*: V_T vs. $V_T - K_s$ colour-magnitude diagrams.

limits of the strength of the Na I doublet $\lambda\lambda$8183.3,8184.8 Å, or *faint* X-ray counterparts with a large uncertainty ellipse. They are tabulated in Table A.11. Among them, we have not accounted for Kiso A–0904 62, a star-like source with blue colours and very faint Hα emission in only one epoch out of three in Wiramihardja et al. (1989).

We have looked for new cluster member candidates without known features of youth. Since the number of confirmed




Fig. 4. i vs. $i - K_s$ colour-magnitude diagrams of one comparison field. Solid lines are for $i - K_s$ percentiles x = 0.90, 0.97, 0.995, from left to right.

young objects in the analysed regions (58) is relatively low, we cannot define a lower envelope of association members as Caballero (2008c) did in the σ Orionis cluster (he used 241 young objects). Besides, we also want to avoid the uncertainties at very young ages associated to theoretical models (Baraffe et al. 2003), and choose a selection criterion as conservative, neutral, reproducible, and objective as possible. For the photometric selection, we have used the data of the ~87000 sources in the comparison fields to determine the locations in the i vs. $i - K_s$ colour-magnitude diagrams of the Alnilam and Mintaka fields where the probability of contamination by fore- and background sources is minimum. Figs. 4 and 5 illustrate the selection procedure.

First, we have divided the i vs. $i - K_s$ diagram (Fig. 4) of all the comparison fields in eleven horizontal strips of width Δi = 0.5 mag between i = 13.0 and 18.5 mag, and two wider strips between i = 10.0 and 13.0 mag. Since the DENIS catalogue fails to provide accurate photometry for the brightest stars (due to saturation and non-linear effects in their detectors), we will only investigate sources with $i > 10.0$ mag in this section. Secondly, we have computed for each strip the $i - K_s$ colour of the source that lefts redwards of it the $100(1 - x)$ % of the remaining objects, where $x \lesssim 1$ (e.g. the percentiles x = 0.90, 0.97, 0.995 separate the 10, 3 and 0.5 % reddest objects, respectively). For a colour-magnitude diagram and a fixed value of x, there are 13 different values of $i - K_s$, one for each strip. The collection of the 13 values determines a boundary for the selection of association member candidates. Thirdly, we have counted the number of objects redder than the selection boundary for different values of the percentile x. We plot in Figs. 4 and 5 the boundaries for x = 0.90, 0.97, 0.995 in the comparison colour-magnitude diagram, and only for x = 0.995 in the Alnilam and Mintaka diagrams. This is the value actually used for the selection. The percentile x = 0.995 maximises the ratio between the number of objects redder than the boundary in the Orion Belt fields and the number of expected contaminants. As a first order approximation, there should be about 44 objects redder than the x = 0.995 boundary in each of the Alnilam and Mintaka fields ($\sim 87000\frac{1}{10}0.005 \approx 44$). The actual figures of sources redder than the x = 0.995 boundary are 272 and 157 in the Alnilam and Mintaka fields, respectively. Accounting for the incomplete coverage of the DENIS survey (see Section 3.1), we estimate average frequencies of contamination at ~25 and ~33 % for the two Orion Belt fields. Using larger (smaller) values of x would lead to lower (larger) frequencies of contamination, but also to smaller (larger) number of photometric association member candidates.

The frequencies of contamination actually are lower, since many of the sources to the red of the selection criterion are at different heliocentric distances to the Ori OB1b association. We have complemented our DENIS/2MASS data with information in the literature, astro-photometric data from the USNO-B2.0 catalogue, and visual inspection of digitized phtographic plates. The 429 sources have been investigated, one by one, to ascertain their membership in association. Eventually, we classify 167 of them as DENIS/2MASS fore- and background sources in the Alnilam and Mintaka areas based on different criteria (optical/near-infrared colours, proper motions, location in a cometary globule, extended point spread functions):

- Table A.12 shows three intermediate and late F-type stars in the foreground, one nearby high-proper motion star (G 99–18), one distant Mira Cet variable star (X Ori) and two previously unknown sources with very red colours (Ruber 1 and 2). While the former five stars were already known to contaminate the Orion field, the latter two stars are identified here for the first time. One of them (Ruber 1) is a very late M dwarf in the foreground, with an appreciable proper motion, while the other star (Ruber 2) seems to be a pulsating giant in the distant Berkeley 20 open cluster (see details in the notes to Table A.12).
- Seven probable reddened sources in the direction of two dense cometary globulae are listed in Table A.13. Six of them fall in the direction of the IC 423 Bok globule. This molecular cloud harbours the T Tauri star IRAS 05307–0038 and has been also classified as a reflection nebula ([RK68] 29; see note on IRAS 05307–0038 in Table A.10). The globule and the corresponding reddened sources are shown in Fig. A.1. The remaining probable reddened source lies close to the centre of the Ori I–2 globule. There is not enough information to determine whether the seven sources are reddened background stars or very young (Class I/II-like) objects embedded in the globules. A spectroscopic follow-up is necessary to ascertain their actual status.
- Tables A.14 and A.15 provide the 2MASS/2MASX designations of 152 galaxies. The vast majority of them appear tabulated in the Two-Micron All Sky Survey Catalog of Extended Sources, 2MASX (Jarret et al. 2000), and are, therefore, catalogued in the NASA/IPAC Extragalactic Database (NED). Some of them also appeared in the works by Paturel et al. (1989) and Monnier Ragaigne et al. (2003) or in radio catalogues (see notes on PMN J0534–0044 in Table A.15). The red colours of many galaxies can be ascribed to their intrinsic nature (starsbursts, ellipticals,




Fig. 5. Same as Fig. 4, but for the Alnilam (left) and Mintaka (right) fields. Solid lines are for $i - K_s$ percentile $x = 0.995$. Objects to the red of this line are marked with (red) filled circles.

bulges of spirals, and mergers). The extragalactic radio source complex 4C–01.06 could not be identified by us.

– Finally, there is one additional red DENIS/2MASS source with poor photometry, 2MASS J05380010–0122377. It was rejected during a visual inspection: it is a binary object partially resolved in the Digital Sky Survey images that probably does not belong to the Ori OB1b association.

Accounting for the known young stars in the association, foreground dwarfs, background giants, reddened stars and galaxies in Tables A.9 to A.15 that are redwards of the $i - K_s$ percentile $x = 0.995$, there remain 189 and 102 photometric association member candidates in the Alnilam and Mintaka fields, respectively (all published stars of unknown status – Table A.11– are bluewards of the selection criterion). The 291 sources are tabulated in Tables A.16 and A.17[5].

Ten of them were photometric member candidates of the "ϵ Orionis cluster" in Scholz & Eislöffel (2005). They classified three of these sources, with identification numbers 44, 120 and 126, as candidates with significant periodic variability (see details in notes to Tables A.16 and A.17). V993 Ori is also a bright photometric variable (V993 Ori; Luyten 1932). Photometric variability is a very common feature in young stars in general and T Tauri stars in particular (e.g. Bertout 1989). There are also variable T Tauri substellar analogs (Caballero et al. 2006). The 280 remaining red DENIS/2MASS sources are firstly shown in our work. We use the acronyms "Annizam" and "Mantaqah" plus running numbers for naming the objects in the Alnilam and Mintaka fields, respectively[6]. The running numbers indicate the position of the association members and candidates with respect to the two supergiants. The three last digits are for the position angle, while the three or four first digits are for the angular separation (for example, Annizam 1751268 is located at $\rho \approx 1751$ arcsec [20.8 arcmin] and $\theta \approx 268$ arcsec with respect to Alnilam). This designation is similar to the Mayrit nomenclature for σ Orionis cluster members and candidates (Caballero 2008c).

2.4. Remarkable fore- and background objects

For completeness, in Table A.18 we list four very bright nearby stars ($K_s \leq 4.5$ mag; 19 Lep, HD 43429, θ Lep, η Lep), a recently-identified very bright He-B subdwarf (Albus 1 – Caballero & Solano 2007; Vennes, Kawka & Allyn Smith 2007), two unknown Tycho-2 high proper motion stars with $\mu > 120$ mas a^{-1}, and six optical counterparts of *IRAS* sources with very red colours ($V_T - K_s > 5.4$ mag) which fall in the comparison fields. Only one of the *IRAS* sources had previously been investigated in the literature, CSS 205, which was classified as a S-type star by Stephenson (1984). Some of the other five stars are even redder than CSS 205, indicating that they could be C- or S-type stars as well. Their very red colours and strong mid-infrared flux excess indicate that they might be stars in the last stages of the AGB phase, or close to the post-AGB stage and evolving into the planetary nebula phase (van der Veen, Habing & Geballe 1989; Trams et al. 1991; Riera et al. 1995).

3. Discussion

Accounting for stars in the tiny overlapping region between the Alnilam and Mintaka fields, we have identified 78 bright early-type (Tables A.3 and A.4) and 58 intermediate and late-type stars (Tables A.9 and A.10) with signatures of youth. Many of the 65 Tycho-2/2MASS (Tables A.5 and A.6) and 17 DENIS/2MASS (Table A.11) published stars of unknown association membership status may also be young. Together with the 291 DENIS/2MASS photometric member candidates

[5] Since there are known young stars in the association and foreground dwarfs bluewards of the selection criterion, and photometric association member candidates in the overlapping region between survey areas, the count of DENIS/MASS sources does not *seem* to coincide: 50 + 167 + 291 ≠ 429.

[6] The name Alnilam derives from the Arabic *an-niż* $\bar{a}m$, related to the word *naẓm*, "string of pearls". The name Mintaka comes from the Arabic *manṭaqah*, "belt".




Fig. 6. Same as Fig. 2, but for the DENIS/2MASS sources. Note the missing DENIS strips.

of the Ori OB1b association (Tables A.16 and A.17), this makes a catalogue of 509 confirmed and candidate young stars and brown dwarfs.

For the canonical age of 5 Ma for the Ori OB1b association, the heliocentric distance of $d = 388\pm30$ pc to the spectroscopic eclipsing binary VV Ori (Terrell et al. 2007) and the colour excess of the supergiant Alnilam $E(B-V) = 0.09$ mag (Lee 1968), and using the DUSTY00 models of the Lyon group (Chabrier et al. 2000), we estimate that the star-brown dwarf boundary (at $M \approx 0.072\,M_{\odot}$ for solar metallicity) in the region is at $J \approx 15.5\pm0.2$ mag. The estimation is identical if the NEXTGEN98 models are used (Baraffe et al. 1998). The objects [SE2005] 126 and Mantaqah 2691223, which are fainter than this magnitude, are the only candidate young brown dwarfs in our work. The star-brown dwarf boundary in the Alnilam-Mintaka region is ∼1 mag fainter than in the σ Orionis cluster (at about $J \sim 14.5$ mag; Caballero et al. 2007), which is younger and *supposed to be* slightly closer. A different heliocentric distance to Ori OB1b, d' (e.g. 330 pc, 440 pc; see Sherry [2003]), would simply shift the star-brown dwarf boundary by a factor $\Delta J = 5 \log d/d'$ (about 0.3 mag in the examples above), which is of the order of the uncertainty in the magnitude limit. Since we have *not* used the heliocentric distance in any step of the association member selection, a different d' would only affect the actual number of substellar objects in our catalogue (the closest distance would lead to have 11 and 5 brown dwarfs in the Alnilam and Mintaka regions, respectively; there would be no brown dwarfs for the farthest distance).

Apart from being 20–40 % more massive than σ Orionis members of the same apparent magnitude, the young objects in the Alnilam-Mintaka region are also distributed along a wider area. Caballero (2008c) identified 75 very low-mass stars, brown dwarfs, and candidates fainter than $J = 14.0$ mag from a DENIS/2MASS correlation, very similar to that presented here, but in a smaller area (a circle of radius 30 arcmin) centred on the Trapezium-like σ Ori system. Accounting for the factor 2.25 of the different survey areas ($\pi 45^2/\pi 30^2$), we expected to have found within the completeness ∼170 young objects with $J > 14.0$ mag in each Orion Belt field *if* the surface densities there and in σ Orionis were identical. Actually, a total of 131 and 63 intermediate- and late-type photometric member candidates of the Ori OB1b association in Tables A.16 and A.17 are fainter than $J = 14.0$ mag (i.e. $M \lesssim 0.09\,M_{\odot}$). We have identified, therefore, ∼70 and ∼40 % less of the expected number of low-mass stars in the Alnilam and Mintaka fields, respectively. Assuming also a similarity in mass functions, it is deduced, therefore, that the surface density of brown dwarfs in Alnilam-Mintaka should be ∼70–40 % of that in σ Orionis. In this computation, we have not taken into account the different location of the star-brown dwarf boundaries. Far from being pessimistic, the lower (sub)stellar density surrounding the two supergiants suggests to survey only ∼1.3 (Alnilam) and ∼2.7 (Mintaka) times more area to find the same number of brown dwarfs than in σ Orionis. A coarse extrapolation of the number of brown dwarf candidates and possible contaminants in Béjar et al. (2003b) and Scholz & Eislöffel (2005) supports our estimations of a relatively high substellar surface density surrounding Alnilam.

The depth of the DENIS survey ($i_{5\sigma} \sim 18.0$ mag) and the expected red colours of young brown dwarfs in the Ori OB 1b association ($i - J \geq 2.5$ mag) has allowed our search to be complete only down to $M \sim 0.08$–$0.07\,M_{\odot}$ ($M \sim 0.05$ and $0.10\,M_{\odot}$ for $d' = 330$ and 440 pc, respectively). Besides, the spatial coverage is incomplete (some strips of the DENIS survey are absent; see Fig. 6) and an important fraction of the actual intermediate- and late-type members of the association may lie bluewards of the conservative $i - K_s$ selection criterion used in Section 2.3. Even accounting for these incompletenesses and for possible contamination among the photometric association member candidates, our work is by far the most comprehensive star compilation in the Alnilam-Minataka region.

Our compilation is not only useful for probing the stellar and substellar populations in the Alnilam-Mintaka region, but also for investigating the mass function in the whole stellar domain from ∼15 to ∼ 0.08–0.07 $M_{\odot}$, the spatial distribution, or the frequency of discs (there are a large amount of association members and candidates with *IRAS* flux excess and/or red near-infrared colours, $J - K_s > 1.15$ mag). Some of these works will be carried out in the near future (Caballero & Solano, in prep.). In the next Section, we present a preliminar study of the radial

distribution of young stars and candidates surrounding Alnilam and Mintaka.

3.1. Spatial distribution

In total, we catalogue 89 confirmed stars and 189 DENIS/2MASS photometric association member candidates in the Alnilam field; in the Mintaka field, we catalogue 47 confirmed stars and 102 DENIS/2MASS photometric association member candidates. The spatial distribution of both type of objects, displayed in top panels in Fig. 7, shows no *clear* radial concentration towards the supergiants. Following the σ Orionis cluster parallelism, we expected a radial density gradient centred on the massive OB-type stars.

We have investigated the normalized cumulative number of confirmed stars and photometric association member candidates in projection within a distance r to Alnilam and Mintaka, $f(r)$. Here, we follow the procedure detailed by Caballero (2008a). Given the incomplete coverage of DENIS in the survey fields (evident in Fig. 6), and to maintain the *radial* symmetry, we have only accounted for the association member candidates to the east of the supergiants. The radial distributions for both confirmed and candidate young objects, shown in the bottom panels of Fig. 7, resemble each other within Poissonian errorbars (not shown for clarity). This resemblance suggests that there is no clear bias in our data compilation (e.g. there has not been a tendency in the literature to survey close to the central OB-type stars, like in σ Orionis). In the bottom panels, we also plot three theoretical power-law distributions. The distribution $f(r) \propto r^2$ corresponds to a uniform distribution of objects in the survey area. The observed radial distribution in σ Orionis is much more radially concentrated, with $f(r) \propto r^1$ in the innermost 20 arcmin-radius cluster core. This distribution corresponds to a volume density proportional to r^{-2}, which is consistent with the collapse of an isothermal spherical molecular cloud (see again details in Caballero [2008a]). Next, we show our results for the Alnilam and Mintaka regions.

3.1.1. Alnilam

Within the uncertainties and the incompleteness of our catalogue, the distribution of young stars and candidates surrounding Alnilam clearly departs from a radially concentrated distribution, as found in σ Orionis. The distribution at less than 25 arcmin to Alnilam is fairly fit by a uniform spacing, while there is a hint of an overdensity of young stars at larger separations. This result is in accord with the classical view of Collinder 70 (see Section 1.1) being a sparse, very wide clustering that might extend to, and overlap with, neighbouring regions (e.g. Mintaka or the "halo" of the σ Orionis cluster – Caballero 2008a). Clues of a σ Orionis-like cluster around Alnilam were not found either by Sherry (2003); his survey had, however, a less extensive spatial coverage, which would increase the difficulty in identifying a "weak cluster".

It is important to notice the great difference between the "cores" of Collinder 70 and σ Orionis. While there are ~130 known stars and brown dwarfs in the innermost 10 arcmin of the latter cluster (most of them with features of youth), we estimate that there are no more than 50 stars in the same area centred on Alnilam. This is not an observational bias, because this abrupt deficiency is not detected in the Mintaka field (Alnilam and Mintaka have roughly the same magnitudes, and so do the sizes of their optical glares – an intense background by a nearby bright star may prevent the detection of sources within the completeness in a photometric survey). However, the frequency of young stars and brown dwarfs at intermediate separations from Alnilam (e.g. 25–45 arcmin) can be larger than in the same corona centred on σ Ori. In any case, a wider study of the radial distribution of young stars, covering the whole Orion Belt, is needed to ascertain the real nature of the Collinder 70 cluster.

3.1.2. Mintaka

The radial distribution of young stars surrounding Mintaka follows, in contrast to Collinder 70, a power-law with an odd exponent intermediate between 1 and 2. This radial concentration may suggest that there is actually a clustering of young stars surrounding the supergiant. This is the first time to propose the existence of a cluster in the area, to which we call "Mintaka cluster". A radial distribution with a power-law $f(r) \propto r^{1.5}$ would correspond to a volume density proportional to $r^{-1.5}$. The lower central concentration than in the scenario of collapse of an isothermal spherical molecular cloud may suggest that (*i*) the Mintaka cluster formed from a non-isothermal molecular cloud, (*ii*) it formed from an isothermal molecular cloud but next suffered from dynamical evolution (and the Mintaka cluster would be, in this scenario, a dynamically-evolved analog to σ Orionis), or (*iii*) there is significant overlapping between the stellar population of Ori OB1b and Ori OB1a. The Ori OB1a sub-association is expected to have a spatial distribution that completely overlaps with the region near Mintaka. Ori OB1a is older than Ori OB1b, but also up to ~100 pc nearer (Sherry 2003). As a result, the isochrones for Ori OB1a and b roughly fall on the same location in the colour-magnitude diagram. Stars from Ori OB1a would have indicators of youth as well, but would not follow the clustering around Mintaka. They would also be difficult to disentangle from proper motions due to the unfavourable direction of Orion and the Sun's relative motions. The combination of a clustered population (the "Mintaka cluster" in Ori OB1b) with a nearly uniformly distributed population (Ori OB1a) would take an $f(r)$ spatial distribution intermediate between r^1 and r^2. Obviously, further work is required to derive any conclusion.

4. Summary

In search for new hunting grounds for substellar objects, we have investigated the stellar populations surrounding two bright supergiants in the Ori OB 1b association (the Orion Belt). The two very young supergiants are Alnilam (ϵ Ori) and Mintaka (δ Ori). Accounting for all the Tycho-2, DENIS and 2MASS sources in the twelve 45 arcmin-radius main and comparison fields, we have examined 107 434 sources in total. After a comprehensive, inclusive, massive Virtual Observatory analysis and bibliographic data compilation, we list:




Fig. 7. Radial distribution of young stars and brown dwarfs surrounding Alnilam (left) and Mintaka (right). *Top panels:* spatial distribution of confirmed young stars (–red– open stars) and photometric candidates to the east of the supergiants (–blue– dots). *Bottom panels:* normalized cumulative number, $f(r)$, of confirmed young stars (–red– thick solid line) and photometric candidates (–blue– dotted line). The –black– dashed lines indicate the theoretical power-law distributions for $f(r) \propto r^{1/2}$, r^1, r^2 (from top to bottom). Compare with figs. 3 and 4 in Caballero (2008a).

– 78 bright (Tycho-2/2MASS) stars with features of extreme youth (i.e. with very early spectral types or HAeBe signatures),
– 58 intermediate- and late-type (DENIS/2MASS) stars with features of extreme youth (i.e. with Li I in absorption, Hα in emission, low g spectroscopic signatures, X-ray emission),
– 289 intermediate- and late-type (DENIS/2MASS) young star candidates with very red $i - K_s$ colours,
– 2 brown dwarf candidates with very red $i - K_s$ colours: [SE2005] 126 (Mantaqah 1582164) and Mantaqah 2691223 (their actual substellar nature depends on the heliocentric distances and ages),
– 82 (Tycho-2/2MASS and DENIS/2MASS) stars without clear signposts of youth that might also belong to the Ori OB 1b association.
– 117 (Tycho-2/2MASS and DENIS/2MASS) stars in the fore- or the background based on their proper motions, spectral types, parallactic heliocentric distances, radial velocities and/or colours,
– 152 extended galaxies, and
– 13 remarkable fore- and backgound stars in the comparison fields.

We report for the first time X-ray emission, *IRAS* flux excess and possible resolved multiplicity for dozens young stars and candidates in the Alnilam-Mintaka region. This abundance represents an excellent compilation of candidates for further follow-up dedicated studies. The vast majority of the listed association member candidates are new. A wealth of detailed information is provided in Appendix A for about one hundred investigated sources.

Finally, we investigate the spatial distribution of stars surrounding Alnilam and Mintaka, and discuss on the possibilities for searching for brown dwarfs. Collinder 70, the cluster that surrounds Alnilam, *if* it exists, must be larger than our search radius of 45 arcmin. Its (sub)stellar population may, therefore, spatially overlap with neighbouring star-forming regions, like the σ Orionis cluster, which is one of the richest regions in

substellar objects. The evidence for a real cluster surrounding Mintaka is, however, more apparent but not conclusive from our data analysis. The "Mintaka cluster", that is presented here for the first time, is less concentrated than the σ Orionis cluster and might represent a next evolutionary stage of it. Accounting for the fainter star-brown dwarf boundary and the lower spatial density of stars very close to the supergiants with respect to σ Orionis, the clusters surrounding Alnilam and Mintaka can be considered to be "elder brothers" (in contraposition to "fraternal twins") of σ Orionis.

Acknowledgements. We appreciate the skillfull referee report by W. H. Sherry. J.A.C. formerly was an Alexander von Humboldt Fellow at the MPIA and currently is an Investigador Juan de la Cierva at the UCM. We thank V. J. S. Béjar, F. Fontanot, S. More, R. Mundt, A. Sicilia-Aguilar and S. Wolf for helpful comments. Partial financial support was provided by the Universidad Complutense de Madrid, the Spanish Virtual Observatory and the Spanish Ministerio Educación y Ciencia under grants AyA2005–02750, AyA2005–04286 and AyA2005–24102–E of the Programa Nacional de Astronomía y Astrofísica and by the Comunidad Autónoma de Madrid under grant CCG07–UCM/ESP–2679 and PRICIT project S–0505/ESP–0237 (AstroCAM). This research has made use of: the SIMBAD, operated at Centre de Données astronomiques de Strasbourg, France; TOPCAT (Tool for OPerations on Tables And Catalogues), provided by the AstroGrid Virtual Observatory project; the NASA's Astrophysics Data System as bibliographic service; and the NASA/IPAC Extragalactic Database (NED) which is operated by the Jet Propulsion Laboratory, California Institute of Technology, under contract with NASA.

References

Abbott, D. C., Bieging, J. H., Churchwell, E. & Cassinelli, J. P. 1980, ApJ, 242, 1183
Abt, H. A. 1979, ApJ, 230, 485
Abt, H. A. & Levato, H. 1977, PASP, 89, 797
Adams, W. S. 1912, ApJ, 35, 163
Aitken, R. G. & Doolittle, E. 1932, New general catalogue of double stars within 120 deg of the North pole..., Carnegie institution of Washington, Washington, D.C. (1932)
Alcalá, J. M., Terranegra, L., Wichmann, R., Chavarría-K.,C., Krautter, J., Schmitt, J. H. M. M., Moreno-Corral, M. A., de Lara E. & Wagner, R. M. 1996, A&AS, 119, 7
Alcalá, J. M., Covino, E., Torres, G., Sterzik, M. F., Pfeiffer, M. J. & Neuhuser, R. 2000, A&A, 353, 186
Álvarez, C., Hoare, M., Glindemann, A. & Richichi, A. 2004, A&A, 427, 505
Andrillat, Y. & Jaschek, C. 1998, A&A, 337, 512
Anthony-Twarog, B. J. 1982, AJ, 87, 1213
Artyukhina, N. M. & Karimova, D. K. 1959, SvA, 3, 122
Bagnulo, S., Landstreet, J. D., Mason, E., Andretta, V., Silaj, J. & Wade, G. A. 2006, A&A, 450, 777
Baraffe, I., Chabrier, G., Barman, T. S., Allard, F. & Hauschildt, P. H. 2003, A&A, 402, 701
Barbaro, G., Mazzei, P., Morbidelli, L., Patriarchi, P. & Perinotto, M. 2001, A&A, 365, 157
Bartaya, R. 1974, ATsir, 845, 3
Bate, M. R., Bonnell, I. A. & Bromm, V. 2003, MNRAS, 339, 577
Becker, R. H., White, R. L. & Edwards, A. L. 1991, ApJS, 75, 1
Béjar, V. J. S. 2001, PhD thesis, Universidad de La Laguna, Spain
Béjar, V. J. S., Zapatero Osorio, M. R. & Rebolo, R. 1999, ApJ, 521, 671
Béjar, V. J. S., Rebolo, R., Zapatero Osorio, M. R. & Caballero, J. A. 2003a, The Future of Cool-Star Astrophysics: 12th Cambridge Workshop on Cool Stars, Stellar Systems, and the Sun (2001 July 30 – August 3), eds. A. Brown, G. M. Harper, and T. R. Ayres, (University of Colorado), 2003, p. 651–657
Béjar, V. J. S., Caballero, J. A. & Rebolo, R. 2003b, unpublished contribution at Science with the GTC 10-m telescope, Granada, Spain, 5–8 February 2002
Berghoefer, T. W., Baade, D., Schmitt, J. H. M. M., Kudritzki, R.-P., Puls, J., Hillier, D. J. & Pauldrach, A. W. A. 1996, A&A, 306, 899
Bernacca, P. L. & Ciatti, F. 1972, A&A, 19, 482
Bertout, C. 1989, ARA&A, 27, 351
Bhatt, H. C. & Manoj, P. 2000, A&A, 362, 978
Bidelman, P. W. 1965, PASP, 77, 388
van Biesbroeck, G. 1974, ApJS, 28, 413
Blaauw, A. 1964, ARA&A, 2, 213
Blaauw, A. 1991, The Physics of Star Formation and Early Stellar Evolution, NATO Advanced Science Institutes (ASI) Series C, Vol. 342, held in Agia Pelagia, Crete, Greece, May 27 – June 8, Dordrecht: Kluwer, 1991, edited by Charles J. Lada and Nikolaos D. Kylafis, p.125
Blaauw, A. & van Albada, T. S. 1963, ApJ, 137, 791
Blomme, R., Prinja, R. K., Runacres, M. C. & Colley, S. 2002, A&A, 382, 921
Bohlender, D. A. 1989, ApJ, 346, 459
Bohlender, D. A., Landstreet, J. D., Brown, D. N. & Thompson, I. B. 1987, ApJ, 323, 325
Bohlin, R. C., Savage, B. D. & Drake, J. F. 1978, ApJ, 224, 132
Bois, B., Lanning, H. H. & Mochnacki, S. W. 1988, AJ, 96, 157
Bonnarel, F., Fernique, P., Bienaymé, O., Egret, D., Genova, F., Louys, M., Ochsenbein, F., Wenger, M. & Bartlett, J. G. 2000, A&AS, 143, 33
Bopp, B. W. 1988, AJ, 95, 1543
Borra, E. F. 1981, ApJ, 249, L39
Briceño, C., Hartmann, L., Stauffer, J. & Martín, E. L. 1998, AJ, 115, 2074
Brown, D. N. & Shore, S. N. 1986, NIA, 86, 365
Brown, A. G. A., de Geus, E. J. & de Zeeuw, P. T. 1994, A&A, 289, 101
Burnham, S. W. 1879, Double star observations made in 1877–8 at Chicago with the 18 1/2-inch refractor of the Dearborn observatory, London (1879)
Burnham, S. W. 1906, A general catalogue of double stars within 121 deg of the North pole, Carnegie institution of Washington, University of Chicago press, Chicago (1906)
Bychkov, V. D., Bychkova, L. V. & Madej, J. 2005, A&A 430, 1143
Caballero, J. A. 2007a, A&A, 466, 917
Caballero, J. A. 2007b, A&A, 462, L61
Caballero, J. A. 2008a, MNRAS, 383, 375
Caballero, J. A. 2008b, MNRAS, 383, 750
Caballero, J. A. 2008c, A&A, 478, 667
Caballero, J. A. & Solano, E. 2007, ApJ, 665, L151
Caballero, J. A. & Solano, E. 2008, A&A, submitted
Caballero, J. A., Béjar, V. J. S., Rebolo, R. & Zapatero Osorio, M. R. 2004, A&A, 424, 857
Caballero, J. A., Martín, E. L., Zapatero Osorio, M. R., Béjar, V. J. S., Rebolo, R., Pavlenko, Ya. & Wainscoat, R. 2006, A&A, 445, 143
Caballero, J. A., Béjar, V. J. S., Rebolo, R. et al. 2007, A&A, 470, 903
Campbell, W. W. 1894, AstAP, 13, 476
Cannon, A. J. & Pickering, E. C. 1924, Henry Draper (HD) catalog and HD extension, Annals of the Astronomical Observatory, Cambridge: Harvard College, Astronomical Observatory (1924) [VizieR On-line Data Catalog: III/135A]

Catalano, F. A. & Renson, P. 1998, A&AS, 127, 421
Cederblad, S. 1946, Lund Medd. Astron. Obs. Ser. II, 119, 1
Cernicharo, J., Bachiller, R., Duvert, G., González-Alfonso, E. & Gómez-Gonzalez, J. 1992, A&A, 261, 589
Chabrier, G. & Baraffe, I. 2000, ARA&A, 38, 337
Chary, R. & Elbaz, D. 2001, ApJ, 556, 562
Cherrington, E., Jr. 1937, ApJ, 85, 139
Ciatti, F., Bernacca, P. L. & D'Innocenzo, A. 1978, A&A, 69, 171
Codella, C., Palumbo, G. G. C., Pareschi, G., Scappini, F., Caselli, P. & Attolini, M. R. 1995, MNRAS, 276, 57
Cohen, M. & Kuhi, L. V. 1979, ApJS, 41, 743
Collinder, P. 1931, Ann. Obs. Lund, 2, 1
Condon, J. J., Cotton, W. D., Greisen, E. W., Yin, Q. F., Perley, R. A., Taylor, G. B. & Broderick, J. J. 1998, AJ, 115, 1693
Coulson, I. M., Walther, D. M. & Dent, W. R. F. 1998, MNRAS, 296, 934
Couteau, P., 1962, JO, 45, 43
Crawford, D. L. & Barnes, J. V. 1966, AJ, 71, 611
Crowther, P. A., Lennon, D. J. & Walborn, N. R. 2006, A&A, 446, 279
Cruz-González, C., Recillas-Cruz, E., Costero, R., Peimbert, M. & Torres-Peimbert, S. 1974, RMxAA, 1, 211
Daniel, Z. 1916, Publications of the Allegheny Observatory of the University of Pittsburgh, 3, 179
Delfosse, X., Tinney, C. G., Forveille, T. et al. 1997, A&A, 327, L25
Dias, W. S., Lépine, J. R. D. & Alessi, B. S. 2001, A&A, 376, 441
Dommanget, J. & Nys, O. 1994, Com. de l'Observ. Royal de Belgique, 115, 1 (1994)
Dommanget, J. & Nys, O. 2002, VizieR On-line Data Catalog: I/274. Originally published in: Observations et Travaux 54, 5 (2002)
Dong, Y. S. & Hu, J. Y. 1991, AcApS, 11, 172
Dorschner, J. & Guertler, J. 1963, AN, 287, 257
Douglas, J. N., Bash, F. N., Bozyan, F. A., Torrence, G. W. & Wolfe, C. 1996, AJ, 111, 1945
Downes, R. A. & Keyes, C. D. 1988, AJ, 96, 777
Drake, S. A., Abbott, D. C., Bastian, T. S., Bieging, J. H., Churchwell, E., Dulk, G. & Linsky, J. L. 1987, ApJ, 322, 902
Dreyer, J. L. E. 1888, MmRAS, 49, 1
Dreyer, J. L. E. 1895, MmRAS, 51, 185
Duflot, M., Figon, P. & Meyssonnier, N. 1995, A&AS, 114, 269
Ebbets, D. 1982, ApJS, 48, 399
Eggen, O.J. 1985, PASP, 97, 807
Epchtein, N., de Batz, B., Capoani, L. et al. 1997, Msngr, 87, 27
Faraggiana, R., Bonifacio, P., Caffau, E., Gerbaldi, M. & Nonino, M. 2004, A&A, 425, 615
Fedorovich, V. P. 1960, Perem. Zvezdy, 13, 166
Flesch, E. & Hardcastle, M. J. 2004, A&A, 427, 387
Frost, E. B. 1909, ApJ, 29, 233
Franciosini, E., Pallavicini, R. & Sanz-Forcada, J. 2006, A&A, 446, 501
Friedemann, C., Gürtler, J. & Löwe, M. 1996, A&AS, 117, 205
Fujii, T., Nakada, Y. & Parthasarathy, M. 2002, A&A, 385, 884
García, J. R., Cebral, J. R., Scoccimarro, E. R. et al. 1995, A&AS, 109, 201
García-Lario P., Manchado A., Pych W. & Pottasch S. R. 1997, A&AS, 126, 479
Garrison, R. F. 1967, PASP, 79, 433
Garrison, R. F. 1994, RMxAA, 29, 111
Gerbaldi, M., Faraggiana, R. & Balin, N. 2001, A&A, 379, 162
Gerbaldi, M., Faraggiana, R. & Lai, O. 2003, A&A, 412, 447
de Geus, E. J., Lub, J. & van de Grift, E. 1990, A&AS, 85, 915
Giclas, H. L., Burnham, R. & Thomas, N. G. 1961, LowOB, 5, 61
Gieseking, F. 1983, A&A, 118, 102
Goldman, B., Delfosse, X., Forveille, T. et al. 1999, A&A, 351, L5
Gómez, J. F., de Gregorio-Monsalvo, I., Suárez, O. & Kuiper, T. B. H. 2006, AJ, 132, 1322
González-García, B. M., Zapatero Osorio, M. R., Béjar, V. J. S., Bihain, G., Barrado y Navascués, D., Caballero, J. A. & Morales-Calderón, M. 2006, A&A, 460, 799
Goy, G. 1973, A&AS, 12, 277
Gray, R. O. & Corbally, C. J. 1993, AJ, 106, 632
Grenier, S., Baylac, M.-O., Rolland, L. et al. 1999, A&AS, 137, 451
Griffin, R. F. 1972, MNRAS, 155, 449
Griffith, M. R., Wright, A. E., Burke, B. F. & Ekers, R. D. 1995, ApJS, 97, 347
Grillo, F., Sciortino, S., Micela, G., Vaiana, G. S. & Harnden, F. R., Jr. 1992, ApJS, 81, 795
Groenewegen, M. A. T. & Lamers, H. J. G. L. M. 1989, A&AS, 79, 359
Guetter, H. H. 1976, AJ, 81, 537
Guetter, H. H. 1979, AJ, 84, 1846
Guetter, H. H. 1981, AJ, 86, 1057
Hambly, N. C., MacGillivray, H. T., Read, M. A. et al. 2001, MNRAS, 326, 1279
Hanbury Brown, R., Davis, J. & Allen, L. R. 1974, MNRAS, 167, 121
Handler, G. 1999, IBVS, 4817, 1
Hardie, R. H., Heiser, A. M. & Tolbert, C. R. 1964, ApJ, 140, 1472
Haro, G. & Moreno, A. 1953, BOTT, 1g, 11
Harris, D. E., Forman, W, Gioa, I. M. et al. 1994, *Einstein* Observatory catalog of IPC X-ray sources (1994), SAO HEAD CD-ROM Series I (*Einstein*), Nos. 18–36 [VizieR On-line Data Catalog: IX/13]
Hartmann, J. 1904, ApJ, 19, 268
Harvey, A. S., Stickland, D. J., Howarth, I. D. & Zuiderwijk, E. J. 1987, Obs, 107, 205
Harvin, J. A., Gies, D. R., Bagnuolo, W. G., Penny, L. R. & Thaller, M. L. 2002, ApJ, 565, 1216
Heintz, W. D. 1980, ApJS, 44, 111
Herbig, G. H. & Kameswara Rao, N. 1972, ApJ, 174, 401
Hernández, J., Calvet, N., Hartmann, L., Briceño, C., Sicilia-Aguilar, A. & Berlind, P. 2005, AJ, 129, 856
Hernández, J., Briceño, C., Calvet, N., Hartmann, L., Muzerolle, J. & Quintero, A. 2006, ApJ, 652, 472
Hernández, J., Hartmann, L., Megeath, T. et al. 2007, ApJ, 662, 1067
Hesser, J. E., McClintock, W. & Henry, R. C. 1977, ApJ, 213, 100
Hillenbrand, L. A. & Carpenter, J. M. 2000, ApJ, 540, 236
Hirth, G. A., Mundt, R. & Solf, J. 1997, A&AS, 126, 437
Ho, P. T. P., Martin, R. N. & Barrett, A. H. 1978, ApJ, 221, L117
Høg, E., Kuzmin, A., Bastian, U., Fabricius, C., Kuimov, K., Lindegren, L., Makarov, V. V. & Röser, S. 1998, A&A, 335, L65
Høg, E., Fabricius, C., Makarov, V. V., Urban, S., Corbin, T., Wycoff, G., Bastian, U., Schwekendiek, P. & Wicenec, A. 2000, A&A, 335, L27
Hubble, E. P. 1922, ApJ, 56, 162
Humpreys, R. M. 1978, ApJS, 38, 309
Jaffe, D. T., Zhou, S., Howe, J. E., Herrmann, F., Madden, S. C., Poglitsch, A., van der Werf, P. P., & Stacey, G. J. 1994, ApJ, 436, 203
Jarad, M. M., Hilditch, R. W. & Skillen, I. 1989, MNRAS, 238, 1085
Jarrett, T.-H., Chester, T., Cutri, R., Schneider, S., Rosenberg, J., Huchra, J. P. & Mader, J. 2000, AJ, 120, 298
Jeffers, H. M., van den Bos, W. H., & Greeby, F. M. 1963, Index catalogue of visual double stars, 1961.0, Publications of the Lick Observatory, Mount Hamilton: University of California, Lick Observatory (1963)
Jeffries, R. D., Maxted, P. F. L., Oliveira, J. M. & Naylor, T. 2006, MNRAS, 371, L6

Johnson, H. L. & Morgan, W. W. 1953, ApJ, 117, 313
Joncas, G. & Borra, E. F. 1981, A&A, 94, 134
Jordan, F. C. 1914, PhD theis, University of Chicago
Keeler, J. E. 1894, AstAp, 13, 476
Keenan, P. C. & Hynek, J. A. 1950, ApJ, 111, 1
Kholtygin, A. F., Burlakova, T. E., Fabrika, S. N., Valyavin, G. G. & Yushkin, M. V. 2006, ARep, 50, 887
Kirkpatrick, J. D., Reid, I. N., Liebert, J. et al. 1999, ApJ, 519, 802
Klemola, A. R., Jones, B. F. & Hanson, R. B. 1987, AJ, 94, 501
Koch, R. H. & Hrivnak, B. J. 1981, ApJ, 248, 249
Koen, C. & Eyer, L. 2002, MNRAS, 331, 45
Kogure, T., Yoshida, S., Wiramihardja, S. D., Nakano, M., Iwata, T. & Ogura, K. 1989, PASJ, 41, 1195
Kraemer, K. E., Shipman, R. F., Price, S. D., Mizuno, D. R., Kuchar, T., Carey & S. J. 2003, AJ, 126, 1423
Kramer, C., Stutzki, J. & Winnewisser, G. 1996, A&A, 307, 915
Kudryavtsev, D. O., Romanyuk, I. I., Elkin, V. G. & Paunzen, E. 2006, MNRAS, 372, 1804
Kukarkin, B. V., Kholopov, P. N., Artiukhina, N. M. et al. 1981, Catalogue of suspected variable stars. Moscow, 1951. Acad. of Sciences USSR
Lacy, J. H., Knacke, R., Geballe, T. R. & Tokunaga, A. T. 1994, ApJ, 428, L69
Lamers, H. J. G. L. M. 1972, A&A, 17, 34
Lamers, H. J. G. L. M. 1974, A&A, 37, 237
Lampens, P. & Rufener, F. 1990, A&AS, 83, 145
Larionov, M. G., Parijskij, Y. N., Zhuravlev, V. I., Sidorenkov, V. N., Berlin, A. B. & Nizhel'skii, N. A. 1994, A&AS, 106, 119
Lee, T. A. 1968, ApJ, 152, 913
Lindroos, K. P. 1985, A&AS, 60, 183
Luhman, K. L., Liebert, J. & Rieke, G. H. 1997, ApJ, 489, L165
Luhman, K. L., Rieke, G. H., Young, E. T., Cotera, A. S., Chen, H., Rieke, M. J., Schneider, G. & Thompson, R. I. 2000, ApJ, 540, 1016
Luyten, W. J. 1932, AN, 245, 211
Lyngå G. 1981, The Catalogue of Open Star Clusters, ADCBu, 1, 90
Lyngå G. 1987, PAICz, 69, 121
Mader, S. L., Zealey, W. J., Parker, Q. A. & Masheder, M. R. W. 1999, MNRAS, 310, 331
MacMinn, D., Phelps, R. L., Janes, K. A. & Friel, E. D. 1994, AJ, 107, 1807
Magakian, T. Y. 2003, A&A, 399, 141
Maheswar, G., Manoj, P. & Bhatt, H. C. 2003, A&A, 402, 963
Mannino, G. & Humblet, J. 1955, AnAp, 18, 237
Marchetti, E., Faraggiana, R. & Bonifacio, P. 2001, A&A, 370, 524
Marilli, E., Frasca, A., Covino, E., Alcalá, J. M., Catalano, S., Fernández, M., Arellano Ferro, A., Rubio-Herrera, E. & Spezzi, L. 2007, A&A, 463, 1081
Markarjan, B. J. 1951, Biurakan Soob, 9,6
Martín, E. L., Magazzù, A. & Rebolo, R. 1992, A&A, 257, 186
McAlister, H. A. & Hendry, E. M. 1982, ApJS, 49, 267
McConell, D. J. 1982, A&AS, 48, 355
McDowell, J. C. 1994, *Einstein* Observatory Unscreened IPC Data Archive (1994), SAO HEAD CD-ROM Series I (*Einstein*), Nos. 18–36
McGehee, P. M. 2006, AJ, 131, 2959
Mermilliod, J. C. 1983, A&A, 128, 362
Miller Barr, J. 1904, AN, 24, 145
Miller, N. A., Cassinelli, J. P., Waldron, W. L., MacFarlane, J. J. & Cohen, D. H. 2002, ApJ, 577, 951
Molnar, M. R. 1972, ApJ, 175, 453
Monet, D., Canzian, B., Harris, H., Reid, N., Rhodes, A. & Sell, S. 1998, The PMM USNO-A1.0 Catalogue, VizieR On-line Data Catalog: I/243. Originally published in: US Naval Observatory Flagstaff Station (1997)
Monet, D. G., Levine, S. E., Canzian, B. et al. 2003, AJ, 125, 984
Monnier Ragaigne, D., van Driel, W., Schneider, S. E., Jarrett, T. H. & Balkowski, C. 2003, A&A, 405, 99
Morgan, W. W., Keenan, P. C. & Kellman, E. 1943, An atlas of stellar spectra, with an outline of spectral classification, Chicago (IL), The University of Chicago press (1943)
Morgan, W. W., Abt, H. A. & Tapscott, J. W. 1978, Revised MK Spectral Atlas for stars earlier than the Sun, Williams Bay: Yerkes Observatory, and Tucson: Kitt Peak National Observatory (1978)
Moran, E. C., Helfand, D. J., Becker, R. H. & White, R. L. 1996, ApJ, 461, 127
Morrell, N. & Levato, H. 1991, ApJS, 75, 965
Nakajima, T., Oppenheimer, B. R., Kulkarni, S. R., Golimowski, D. A., Matthews, K. & Durrance, S. T. 1995, Nature, 378, 463
Nakano M., Yamauchi S., Sugitani K., Ogura K. & Kogure T. 1999, PASJ, 51, 1
Nesterov V. V., Kuzmin A. V., Ashimbaeva N. T., Volchkov A. A., Röser S. & Bastian U. 1995, A&AS, 110, 367
Neubauer, F. J. 1936, Lick Observatory Bulletin, 17, 185
Neubauer, F. J. 1943, ApJ, 97, 300
Neuhäuser, R. & Comerón, F. 1998, Science, 282, 83
Nordström, B., Stefanik, R. P., Latham, D. W. & Andersen, J. 1997, A&AS, 126, 21
Nordström, B., Mayor, M., Andersen, J., Holmberg, J., Pont, F., Jørgensen, B. R., Olsen, E. H., Udry, S. & Mowlavi, N. 2004, A&A, 418, 989
North, P. 1984, A&A, 141, 328
Ogura, K. & Sugitani, K. 1998, PASA, 15, 91
Ogura, K., Sugitani, K. & Pickles, A. 2002, AJ, 123, 2597
Oliveira, J. M. & van Loon, J. Th. 2004, A&A, 418, 663
Ochsenbein, F., Bauer, P. & Marcout, J. 2000, A&AS, 143, 221
Oudmaijer, R. D., van der Veen, W. E. C. J., Waters, L. B. F. M., Trams, N. R., Waelkens, C. & Engelsman, E. 1992, A&AS, 96, 625
Padovani, P., Allen, M. G., Rosati, P. & Walton, N. A. 2004, A&A, 424, 545
Palla, F. & Baraffe, I. 2005, A&A, 432, L57
Pallavicini, R., Pasquini, L. & Randich, S. 1992, A&A, 261, 245
Pannekoek, A. 1929, Publ. Astron. Inst. Amsterdam, 2, 63
Paturel, G., Fouquée, P., Bottinelli, L. & Gouguenheim, L. 1989, A&AS, 80, 299
Paunzen, E. 2001, A&A, 373, 633
Paunzen, E. & Gray, R. O 1997, A&AS, 126, 407
Paunzen, E., Handler, G., Weiss, W. W. et al. 2002, A&A, 392, 515
Pedersen, H. & Thomsen, B. 1977, A&AS, 30, 11
Pérez-Garrido, A., Díaz-Sánchez, A. & Villo, I. 2005, AN, 326, 1028
Perryman, M. A. C., Lindegren, L., Kovalevsky, J. et al. 1997, A&A, 323, L49
Pismis, P., Haro, G. & Struve, O. 1950, ApJ, 111, 509
Prinja, R. K., Rivinius, Th., Stahl, O., Kaufer, A., Foing, B. H., Cami, J. & Orlando, S. 2004, A&A, 418, 727
Rebolo, R., Zapatero Osorio, M. R. & Martín, E. L. 1995, Nature, 377, 129
Rebolo, R., Zapatero Osorio, M. R., Madruga, S., Béjar, V. J. S., Arribas, S. & Licandro, J. 1998, Science, 282, 1309
Reipurth, Bo & Clarke, C. 2001, AJ, 122, 432
Reipurth, Bo, Bally, J., Fesen, R. A. & Devine, D. 1998, Nature, 396, 343
Renson, P. 1988, A&AS, 76, 127
Renson, P. 1992, BICDS, 40, 97
Roman, N. G. 1955, ApJS, 2, 195
Rossiter, R. A. 1955, Catalogue of Southern Double Stars, Publications of the Observatory of the University of Michigan, Ann Arbor: University of Michigan (1955)

ROSAT Consortium (*) 2000, ROSAT News, 71 [VizieR On-line Data Catalog: IX/28A] (*: Max-Planck-Institut für extraterrestrische Physik, Goddard Space Flight Center, Smithsonian Astrophysical Observatory, Leicester University, Astrophysikalisches Institut Potsdam)
Rufener, F. & Bartholdi, P. 1982, A&AS, 48, 503
Ruiz, M. T., Leggett, S. K. & Allard, F. 1997, ApJ, 491, L107
Salim, S. & Gould, A. 2003, ApJ, 582, 1011
Sanduleak, N. 1971, PASP, 83, 95
Savage, B. D., Massa, D., Meade, M. & Wesselius, P. R. 1985, ApJS, 59, 397
Schaefer, B. E. 2005, Journal for the History of Astronomy, 36, 1667
Schild, R. E. & Chaffee, F. 1971, ApJ, 169, 529
Schild, R. E. & Cowley, A. P. 1971, A&A, 14, 66
Schmidt, E. G. & Carruthers, G. R. 1993, ApJS, 89, 259
Scholz, A. & Eislöffel, J. 2005, A&A, 429, 1007
Sharpless, S. 1952, ApJ, 116, 251
Sharpless, S. 1959, ApJS, 4, 257
Sharpless, S. 1962, ApJ, 136, 767
Sharpless, S. 1974, AJ, 79, 1073
Sherry, W. H. 2003, Ph.D. thesis, State University New York at Stony Brook
Sherry, W. H., Walter, F. M. & Wolk, S. J. 2000, American Astronomical Society, 197th AAS Meeting, #10.03; Bulletin of the AAS, Vol. 32, p. 1412
Sherry, W. H., Walter, F. M. & Wolk, S. J. 2004, AJ, 128, 2316
Skrutskie, M. F., Cutri, R. M., Stiening, R. et al. 2006, AJ, 131, 1163
Sloan, G. C. & Price, S. D. 1998, ApJS, 119, 141
Speck, A. K., Barlow, M. J., Sylvester, R. J. & Hofmeister, A. M. 2000, A&AS, 146, 437
Stebbins, J. 1915, ApJ, 42, 133
Stephenson, C. B. 1986, ApJ, 300, 779
Stephenson, C. B. & Sanduleak, N. 1977, ApJS, 33, 459
Struve, O. & Luyten, W. J. 1949, ApJ, 110, 160
Subramaniam, A., Gorti, U., Sagar, R. & Bhatt, H. C. 1995, A&A, 302, 86
Templeton, M. R., Mattei, J. A. & Willson, L. A. 2005, AJ, 130, 776
Terrell, D., Munari, U. & Siviero, A. 2007, MNRAS, 374, 530
The, P. S., de Winter, D. & Pérez, M. R. 1994, A&AS, 104, 315
Tsalmantza, P., Kontizas, E., Cambrésy, L., Genova, F., Dapergolas, A. & Kontizas, M. 2006, A&A, 447, 89
Ueda, Y., Ishisaki, Y., Takahashi, T., Makishima, K. & Ohashi, T. 2001, ApJS, 133, 1
Vennes, S., Kawka, A. & Allyn Smith, J. 2007, ApJ, 668, L59
Vieira, S. L. A., Corradi, W. J. B., Alencar, S. H. P., Mendes, L. T. S., Torres, C. A. O., Quast, G. R., Guimarães, M. M. & da Silva, L. 2003, AJ, 126, 2971
Vivas, A. K., Zinn, R., Abad, C. et al. 2004, AJ, 127, 1158
Voges, W., Aschenbach, B., Boller, T. et al. 1999, A&A, 349, 389 [VizieR On-line Data Catalog: IX/10A]
Voges, W., Aschenbach, B., Boller, T. et al. 2000, IAU Circ., 7432, 3. Edited by D. W. E. Green (2000) [VizieR On-line Data Catalog: IX/29]
Waelkens, C., Aerts, C., Kestens, E., Grenon, M. & Eyer, L. 1998, A&A, 330, 215
Walborn, N. R. 1983, ApJ, 268, 195
Walter, F. M., Wolk, S. J., Freyberg, M. & Schmitt, J. H. M. M. 1997, MmSAI, 68, 1081
Warren, W. H., Jr. & Hesser, J. E. 1977, ApJS, 34, 115
Warren, W. H., Jr. & Hesser, J. E. 1978, ApJS, 36, 497
Weaver, W. B. & Jones, G. 1992, ApJS, 78, 239
Weintraub, D. A. 1990, ApJS, 74, 575
White, N. E., Giommi, P. & Angelini, L. 2000. Originally published in: Laboratory for High Energy Astrophysics (LHEA/NASA), Greenbelt (2000) [VizieR On-line Data Catalog: IX/31]
Whitford, A. E. 1958, AJ, 63, 201
Whitworth, A., Bate, M. R., Nordlund, A., Reipurth, Bo & Zinnecker, H. 2007, Protostars and Planets V, B. Reipurth, D. Jewitt, and K. Keil (eds.), University of Arizona Press, Tucson, 2007, p.459–476
Wilson, R. E. 1953, General Catalogue of Stellar Radial Velocities, Carnegie Inst. Washington D.C. Publ. 601,1953 (XXBH) (1953)
Wiramihardja, S. D., Kogure, T., Yoshida, S., Ogura, K. & Nakano, M. 1989, PASJ, 41, 155
Wiramihardja, S. D., Kogure, T., Yoshida, S., Nakano, M., Ogura, K. & Iwata, T. 1991, PASJ, 43, 27
Wolf, M. 1904, 165, 29
Wolk, S. J. 1996, Ph.D. thesis, State University New York at Stony Brook
Woolf, V. M. & Lambert, D. L. 1999, ApJ, 520, L55
Wouterloot, J. G. A. & Walmsley, C. M. 1986, A&A, 168, 237
Wouterloot, J. G. A., Walmsley, C. M. & Henkel, C. 1988, A&A, 203, 367
Wouterloot, J. G. A., Walmsley, C. M. & Henkel, C. 1989, A&A, 215, 131
XMM-Newton Survey Science Centre Consortium 2007. XMM-SSC, Leicester, UK (2007) [VizieR On-line Data Catalog: IX/39]
Yudin, R. V. & Evans, A. 1998, A&AS, 131, 401
Yun, J. L. & Clemens, D.P. 1994, AJ, 108, 612
Yun, J. L., Moreia, M. C., Alves, J.F. & Storm, J. 1997, A&A, 320, 167
Zapatero Osorio, M. R., Béjar, V. J. S., Martín, E. L., Rebolo, R., Barrado y Navascués, D., Bailer-Jones, C. A. L. & Mundt, R. 2000, Science, 290, 103
de Zeeuw, P. T., Hoogerwerf, R., de Bruijne, J. H. J., Brown, A. G. A. & Blaauw, A. 1999, AJ, 117, 354

Appendix A: The Annizam/Mantaqah catalogue

Notes to Table A.3:

* HD 36980 AB is a close binary with $\rho \approx 0.7$ arcsec, $\theta \approx 61$ deg (catalogue of Components of Double and Multiple stars; Dommanget & Nys 1994 – there are no fundamental differences between this edition and the second one [Dommanget & Nys 2002], except for the number of considered sources).
* RX J0535.6–0152 AB is a T Tauri star with a red $V_T - K_s$ colour. It is a G6V-type spectroscopic binary with lithium in absorption (pEW(Li I) = +0.32 Å), partially filled Hα line (pEW(Hα) = +2.40 Å) and X-ray in emission (Alcalá et al. 1996, 2000). RX J0535.6–0152 AB was the third strongest X-ray source in the investigation of 40 weak-line T Tauri stars in Orion by Marilli et al. (2007), with $\log L_X = 30.7^{+0.2}_{-0.7}$ (erg cm^{-2} s^{-1}). These authors found it to be a photometric variable with a period of 1.74 d.
* HD 37285 AB is a visual binary star with $\rho \approx 0.4$ arcsec, $\theta \approx 263$ deg (Dommanget & Nys 1994).
* HD 37389 is embedded in the Ori I–2 Cometary Globule (Ho, Martin & Barrett 1978; Cernicharo et al. 1992; Mader et al. 1999). Oudmaijer et al. (1992) and Coulson, Walther & Dent (1998) have reported infrared and submillimetre flux excesses due to a Vega-like disc. The star has appreciable polarization in the optical (Bhatt & Manoj 2000). Some catalogues tabulate a hypothetical companion, BD–01 985B, at $\rho \sim 5.0$ arcsec, $\theta \sim 350$ deg.
* HD 37149 is a helium-weak star (Bernacca & Ciatti 1972; Renson 1988) with Hα in medium emission (Bidelman 1965). It is likely the UV-emission source [SC93b] 328 (Schmidt & Carruthers 1993).
* HD 290770 was discovered as an emission-line star by Bidelman (1965) and has been classified as a B8–9Ve Herbig Ae/Be star by many other authors (Guetter 1976; Gieseking 1983; Dong & Hu 1991; Nesterov et al. 1995; The et al. 1994). Vieria et al. (2003) found [O I]+[S II] in emission and measured the Hα line in double-peak emission, with the secondary peak having more than half the strength of the primary. Yudin & Evans (1998) found negligible polarization in the optical. Previously unnoticed, HD 290770 has one of the most apparent flux excesses at the *IRAS* passbands in the Ori OB 1 b association (Caballero & Solano, in prep.). Here we report a close visual companion to the star at $\rho \sim 6.8$ arcsec, $\theta \sim 340$ deg, and $\Delta K_s = 4.46 \pm 0.03$ mag fainter. From its $I - J$ and $J - K_s$ colours from DENIS and 2MASS, it seems to be a F–G:-type star in the fore-/background.
* HD 37344 is embedded in the bright-rimmed cloud complex Ori I–2N, close to cloud [OS98] 40C (Ogura & Sugitani 1998).
* HD 290602 is also BD–01 947.
* HD 290674 is also BD–01 977.
* HD 37321 AB is a well-known helium-weak star with a high rotational velocity ($v \sin i \approx 100$ km s^{-1}; Mermilliod 1983) and spectrum variability (Molnar 1972; Garrison 1994 – but see Pedersen & Thomsen 1977). Blaauw & van Albada (1963) proposed the star to be a long-period spectroscopic binary; Morrell & Levato (1991) measured, however, a constant radial velocity of 24 ± 6 km s^{-1} during their monitoring. It is a close binary with $\rho = 0.756 \pm 0.002$ arcsec, $\theta = 14 \pm 1$ deg ($\Delta H_p = 1.62 \pm 0.01$ mag; Perryman et al. 1987). It was only resolved by Tycho-2. We accounted for the $B_T V_T$ magnitude of the A component and the near infrared K_s magnitude of both A and B components as a single object. The system could have a faint, red, third component ($H = 10.98 \pm 0.02$ mag), at $\rho \approx 17.7$ arcsec, $\theta \approx 32$ deg. HD 37321 AB may also be the far-ultraviolet emission source [SC93b] 341 (Schmidt & Carruthers 1993).
* HD 36955 is a peculiar magnetic star with abnormal abundances of Si, Cr and Eu (Gray & Corbally 1993; Kudryavtsev et al. 2006).
* V1247 Ori is a Herbig He/Be star (García-Lario et al. 1997; Fujii et al. 2002) whose non-banded Hα emission was found by McConell (1982). Spectral types from A5III, through A7, to F0V have been provided (Schild & Cowley 1971; Nesterov et al. 1995; Vieira et al. 2003). Vieira et al. (2003) found no forbidden lines in its optical spectrum, but identified an Hα symmetric profile without, or with only very shallow, absorption features. No H_2O, NH_3 or CO radio lines were found by Wouterloot et al. (1986, 1988, 1989). V1247 Ori is, besides, a well-studied δ Scuti star, with $P = 0.096967$ d and peak-to-peak amplitude in the V band of 0.050 mag (Lampens & Rufener 1990; García et al. 1995; Handler 1999). Its SED shows clear excesses from the J band to 60–100 μ (Caballero & Solano, in prep.), and is composed of two components, one warm (1.2–2.2 μm) and other cool (12–100 μm).
* Alnilam (ϵ Ori, 46 Ori, HD 37128; $V = 1.70$ mag) is one of the brightest supergiants in the sky and, therefore, one of the best known stars. The first spectroscopic study was carried out more than a century ago by Campbell (1894) and Keeler (1894). It is a hot, massive, single star in the hydrogen shell burning phase (Lamers 1974; Jarad, Hilditch & Skillen 1989) with photometric and spectroscopic variability (Stebbins 1915; Ebbets 1982; Prinja et al. 2004), Hα, X-ray and radio emission (Cherrington 1937; Abbott et al. 1980; Berghöfer et al. 1996; Blomme et al. 2002), and strong stellar wind and mass loss (Groenewegen & Lamers 1989; Prinja et al. 2001; Crowther, Lennon & Walborn 2006). Alnilam is one of the few early-type stars with determination of the angular diameter using optical interferometry (Hanbury Brown, Davis & Allen 1974). It illuminates the NGC 1990 reflection nebula. Last, it has been used as a bright spectrophotometric standard (B0Ia in the MK classification by Johnson & Morgan 1953), to investigate the interstellar extinction (e.g. Whitford 1958; Bohlin, Savage & Drake 1978) and for comparison with other early-type supergiants (Humphrey 1978). A review of "classic" works on Alnilam can be found in Lamers (1972). The status of "Alnilam B" (BD–01 969B; see Table A.11), at 3 arcmin to the northeast, is unknown.
* HD 37397 is a low-amplitude variable star ($P = 0.572885$ d, $A(V) = 0.00089$ mag; Koen & Eyer 2002) with a constant radial velocity of 22–23 km s^{-1} (Morrell & Levato 1991; Duflot et al. 1995; Grenier et al. 1999).

* VV Ori AB is a double-lined eclipsing binary in a detached configuration (Miller Barr 1904; Adams 1912; Daniel 1916; Struve & Luyten 1949). The two early-type stars, B1.0V+B4.5V, are separated by $13.49\pm0.05\,R_{\odot}$ ($P \approx 1.48$ d; Terrell et al. 2007). The mid-infrared source IRAS 05309–0111 is located at $\rho \sim 16$ arcsec, $\theta \sim 180$ deg, to VV Ori. Friedemann, Gürtler & Löwe (1996) had been the only investigators before us to notice the *IRAS* thermal emission of VV Ori, and attributed it to circumstellar dust. This is very important, because: (*i*) stars with very early spectral types, just as the primary in VV Ori, are not expected to have circumstellar discs at the age of the Ori OB 1 b association, and (*ii*) the disc would surround the binary system (i.e. the inner part of the disc would be at several –tens– astronomical units, while the binary components are separated by ∼0.64 AU). VV Ori may also be associated to the X-ray sources [NYS99] A–01 and 1AXG J053331–0110 (Nakano et al. 1999; Ueda et al. 2001).
* HD 36684 AB is a close binary with $\rho \approx 0.2$ arcsec, $\theta \approx 200$ deg (Dommanget & Nys 1994). It also has a high rotational velocity ($v \sin i \approx 160\,\mathrm{km\,s^{-1}}$; Sharpless 1974).
* HD 290750 is a low-amplitude suspected variable (Rufener & Bartholdi 1982).
* HD 36779 AC forms together with the post-T Tauri star HD 36779 B a likely Lindroos system (Lindroos 1985). HD 36779 AC is, in its turn, a spectroscopic binary (Morrell & Levato 1991).
* HD 37187 might form a new (very wide) Lindroos system together with V583 Ori ($\rho \approx 29.0$ arcsec, $\theta \approx 212$ deg).
* HD 37076 and HD 290671 form the STF 751 double system, with $\rho \approx 15.6$ arcsec, $\theta \approx 124$ deg (Dommanget & Nys 1994). They share Tycho-2 proper motion within the uncertainties. The X-ray emission found by *ROSAT* (with HRI and PSPC) is associated to the faintest component (HD 290671, B9.5V; Caballero et al. in prep.).
* HD 290665 is a SiCrEuSr chemically peculiar star (Bartaya 1974; Schild & Cowley 1971; Guetter 1976; Joncas & Borra 1981; Gieseking 1983). It is also a strong magnetic star ($\langle B_z \rangle \approx -0.17$ T; Bagnulo et al. 2006). Last, HD 290665 has a radial velocity discordant with association membership (Gieseking 1983). It could be, however, a spectroscopic binary.
* V1379 Ori is a slowly pulsating B star (Waelkens et al. 1998).
* HD 290662, a peculiar Vega-like star, was proposed to be a spectroscopic binary by Gieseking (1983) based on low quality data.
* HD 36954 AB is a spectroscopic binary (SB1) with a period $P \sim 4.6$ d (Neubauer 1936; Morrell & Levato 1991).
* HD 37235 is a spectroscopic variable B7–A0:V according to Bernacca & Ciatti (1972). Renson (1992) tabulates it as a He-weak star. The origin of the mid-infrared source IRAS 05344–0044, located at a separation $\rho \sim 35$ arcsec to the south of the star, probably lies on the extended source 2MASX J05365804–0042413, whose spectral energy distribution resembles those of starburst galaxies with large dust content (see, e.g., Chary & Elbaz 2001)
* HD 290648, HD 290660 and HD 290650 are also BD–00 1004, BD–00 1020 and BD–00 1012, respectively.

Notes to Table A.4:

* HD 290515 is at a projected angular separation of 34 arcsec to the background RR Lyr star [VZA2004] 28 (Vivas et al. 2004).
* HD 290492 AB is a close binary with $\rho = 0.739\pm0.005$ arcsec, $\theta = 63.9\pm1.9$ deg ($\Delta b = 0.6\pm0.2$ mag – Rossiter 1955; Marchetti, Faraggiana & Bonifacio 2001). It is a non-variable, mild λ Boo star candidate (Paunzen & Gray 1997; Paunzen et al. 2002 – but see Gerbaldi, Faraggiana & Lai [2003] and Faraggiana et al. [2004]). Paunzen (2001) derived a photometric distance of $d = 279\pm20$ pc, but he erroneously assumed no contamination by the secondary in the spectrum of the primary. HD 290492 AB and the G8III star GSC 04766–02124, which is ∼3 mag fainter in the V band and at ∼24 arcsec to the west, do not share a common proper motion, as has been proposed in the literature.
* SS 28 is a T Tauri star. It has been only investigated by Stephenson & Sanduleak (1977), Bopp (1988), Wiramihardja et al. (1989) and Kogure et al. (1989). It has a double-peaked Hα emission line with intensity about three times that of the continuum, Hβ also in emission and an apparently "filled in" Ca II H and K. According to Bopp (1988), SS 28 resembles some unusual interacting F+B binary systems. It is, besides, an *Einstein* Observatory soft X-ray source. The *ROSAT* Satellite measured afterwards 18 events associated to SS 28 (ROSAT 2000; White et al. 2000). SIMBAD tabulates a quadruple identification (it is also Kiso A–0904 6, Kiso A–0903 234 and 2E 1299).
* BD–00 984 is a chemically peculiar star based on their abnormal abundances of Hg, Mn (Woolf & Lambert 1999 – they classified it as one of the youngest HgMn stars) and Si (Brown & Shore 1986). It forms a curious perfect alignement with two nearby radio sources: [LPZ94] 147 ($\rho \approx 31.5$ arcsec, $\theta \approx 177.0$ deg – $S_\nu(3.9\,\mathrm{GHz}) = 48\pm24$ mJy, Larionov et al. 1994) and TXS 0529–004 ($\rho \approx 1.8$ arcmin, $\theta \approx 178$ deg – $S_\nu(1.4\,\mathrm{GHz}) = 82.2\pm2.5$ mJy, Condon et al. 1998; $S_\nu(0.365\,\mathrm{GHz}) = 214\pm24$ mJy, Douglas et al. 1996). The spectral index of TXS 0529–004 is –0.9, consistent within the uncertainties with thermal Bremsstrahlung emission.
* HD 290500 was classified as a Herbig Ae/Be star by Vieira et al. (2003). They derived A2 spectral type and detected Hα in double-peaked emission, with the secondary peak having less than half the strength of the primary. No forbidden lines were identified. Codella et al. (1995) gave an upper limit for the 22.2 GHz H_2O maser emission of the star. HD 290500 has a mid-infrared flux excess, as measured by *IRAS* (Caballero & Solano, in prep.).
* HD 36841 has been widely used for determining ultraviolet interstellar extinction curves (e.g. Savage et al. 1985; Barbaro et al. 2001). It was formerly considered a late-O-type star (Mannino & Humblet 1955; Goy 1973; Cruz-González et al. 1974).

* V1093 Ori AB (HD 36313) is a variable of α^2 CVn type (North 1984; Catalano & Renson 1998), a helium-weak, silicon, magnetic peculiar (Guetter 1976; Borra 1981; Bychkov, Bychkova & Madej 2005) and a close binary star ($\rho \approx 0.22$ arcsec, $\theta \approx 172.6$ deg – Couteau 1962; van Biesbroeck 1974).
* Mintaka AE–D (δ Ori, 34 Ori, HD 36486; $V = 2.23$ mag) is the most famous star in the Orion belt. It is a very bright triple within a hierarchical quintuple system. Mintaka D (δ Ori Ab) is an early-B-type star at $\rho = 0.267$ arcsec, $\theta = 140$ deg ($\Delta H_P = 1.35 \pm 0.02$ mag), from the tight AE binary (Heintz 1980; McCalister & Hendry 1982; Perryman et al. 1997); it may be a rapid rotator or a spectroscopic binary (Harvin et al. 2002). Mintaka A (O9.5II, δ Ori Aa1) and E (B0.5III, δ Ori Aa2) form an eclipsing spectroscopic binary with a peak-to-peak amplitude $A(H_P) = 0.01$ mag and a period $P = 5.7325$ d (Hartmann 1904; Jordan 1914; Pismis, Haro & Struve 1950; Koch & Hrivnak 1981; Harvey et al. 1987; Harvin et al. 2002; Kholtygin et al. 2006); the binary has suffered from an intense mass loss. The other two components, Mintaka B and C, are described below.
* Mintaka C (δ Ori C, HD 36485) is at $\rho \approx 52.2$ arcsec, $\theta \approx 0.1$ deg to Mintaka AE ($\Delta V_T = 4.417 \pm 0.012$ mag; Høg et al. 2000). It is a helium-strong star (Morgan, Abt & Tapscott 1978; Walborn 1983; Bohlender 1989) with non-thermal radio emission (Drake et al. 1987).
* HD 290491 is also TYC 4766 2264 1.
* HD 36726 A is a λ Boo star (Abt & Levato 1977; Paunzen 2001) and has a quite high rotational velocity, $v \sin i$ (about 120 km s^{-1}; Abt 1979). It is the brightest component of a triple system tabulated by Aitken & Doolittle (1932). The secondary is BD–00 993B (Table A.11). The system could be quadruple, since there is an additional 2MASS source at $\rho \approx 5.7$ arcsec, $\theta \approx 3$ deg to HD 36726 A with $J = 15.109 \pm 0.137$ mag.
* HD 290572 is a B8V and a K0V (sic) according to Cannon & Pickering (1924) and Nesterov et al. (1995), respectively. An intermediate A spectral type better matches the $B_T - K_s$ colour.
* HD 290569 is an A0V according to Nesterov et al. (1995); however, the relatively red colour $B_T - K_s = 1.29 \pm 0.12$ mag better matches with a later spectral type (i.e. intermediate A).

Notes to Table A.5:

* HD 37038 AB is a binary whose components are separated by $\rho \approx 0.6$ arcsec, $\theta \approx 265$ deg. The secondary is 2 mag fainter than the primary (Dommanget & Nys 1994). It could actually be a hierarchical triple, since Nordström et al. (1997) found the F-type dwarf to be a double-lined spectroscopic binary with evident radial-velocity variations in scales of a few days (the resolved binary cannot be responsible of such variations).
* HD 36863 has a radial velocity that deviates more than 25 km s^{-1} with respect to the average radial velocity of the association (Gieseking 1983). It satisfies, however, the photometric and proper motion criteria of very young stars in Orion. HD 36863 might be a very young single-line spectroscopic binary (SB1). Guetter (1976) classified it as an A7-type star.
* [NYS99] A–06 is an X-ray source identified by *ASCA* and *ROSAT* (1AXG J053448–0131, 1RXS J053450.5–013120). We estimate F: spectral type from its $V_T - K_s$ colour.
* HD 290673 is also BD–01 975.
* TYC 4767 1130 1 seems to be a close ($\rho \sim 3$ arcsec) binary from the 2MASS data.
* HD 37172 is a probable non-member of the association according to Guetter (1976), although other authors consider it to be a real member (Warren & Hesser 1977, 1978; Hesser, McClintock & Henry 1977; Gieseking 1983; de Geus & van de Grift 1990). It has a peculiar Mn I λ4030–4035 Å blend (Gray & Corbally 1993) and falls slightly to the red of the association sequence in the $V_T - K_s$ vs. V_T diagram (Fig. 3). It is at $\rho \approx 5.0$ arcmin, $\theta \approx 115$ deg, to Alnilam.
* TYC 4766 2371 1 is the third closest bright star to Alnilam ($\rho \approx 6.0$ arcmin, $\theta \approx 250$ deg).
* TYC 4766 2150 1 (CCDM J05375–0103B) has been repeatedly proposed to form a binary system together with HD 290749 ($\rho \approx 29.0$ arcsec, $\theta \approx 344$ deg – Burnham 1906; Dommanget & Nys 1994). They do not share, however, a common proper motion.
* TYC 4766 542 1 might be the X-ray source 1RXS J053447.9–010224 (1AXG J053446–0102; Ueda et al. 2001). The nearby binary 2MASS J05344642–0102340 ($\rho \approx 4.0$ arcsec, $\theta \approx 220$ deg, $\Delta H = 1.89 \pm 0.05$ mag), at ∼35 arcsec to the west of TYC 4766 542 1 could also be the X-ray source.
* HD 290746 is a G0V according to Nesterov et al. (1995); however, the relatively blue colour $B_T - K_s = 1.38 \pm 0.09$ mag better matches with an earlier spectral type (i.e. late A or early F).

Notes to Table A.6:

* HD 290585 has a debris disc according to MIPS/*Spitzer* observations at 24 μm by Hernández et al. (2006). It is a double binary resolved by 2MASS ($\rho \approx 5.6$ arcsec, $\theta \approx 132$ deg, $\Delta H = 2.05 \pm 0.05$ mag); the secondary is not in the Tycho-2 catalogue.
* HD 290513 is a F0V dwarf according to Nesterov et al. (1995); a G–K spectral type better matches the observed colours.
* TYC 4766 790 1 has a visual companion at $\rho \approx 8.5$ arcsec, $\theta \approx 190$ deg ($\Delta H = 0.69 \pm 0.04$ mag).
* HD 290583 A has a visual companion of roughly the same brightness. HD 290583 B, not identified by Tycho-2, is at $\rho \approx 7.44$ arcsec, $\theta \approx 3.8$ deg ($\Delta H = 0.40 \pm 0.03$ mag).
* HD 290507 and HD 290504 are A5V dwarfs according to Nesterov et al. (1995); F–G spectral types better match the observed colours.
* TYC 4766 528 1 is the brightest component of a visual triple system. The other two components are located at $\rho \approx$

3.4 arcsec, $\theta \approx 165$ deg ($\Delta H = 0.36 \pm 0.07$ mag), and $\rho \approx 7.4$ arcsec, $\theta \approx 195$ deg ($\Delta H = 2.07 \pm 0.06$ mag).
* TYC 4766 1168 1 has a visual companion at $\rho \approx 9.4$ arcsec, $\theta \approx 10$ deg ($\Delta H = 0.98 \pm 0.04$ mag).
* TYC 4766 2424 1 has a low a proper motion of less than 5 mas a^{-1} and a blue colour $B_T - K_s = 0.63 \pm 0.09$ mag, typical of early A-type stars in the association.
* TYC 4753 49 1 is at $\rho \sim 46$ arcsec, $\theta \sim 288$ deg to the radio source [LPZ94] 146 (Larionov et al. 1994).

Notes to Table A.7:

* IRAS 05354–0142 has the reddest $V_T - K_s$ colour among the ∼1500 Tycho-2/2MASS investigated stars in the survey area ($V_T - K_s = 7.1 \pm 0.3$ mag). The closeness of IRAS 05354–0142 to the Ori I–2 globule may explain part of its reddening, but not all. It might be a S-type or a C-type giant with a very late spectral type and very low effective temperature. The absence of a mid-infrared excess (Kraemer et al. 2003) rules out the hypothesis of IRAS 05354–0142 being a proto-star in the upper part of the Hayashi track of collapse associated to the Bok globule.
* HD 290680 has a $V_T - K_s$ colour that better matches with a K spectral type.
* HD 290679 is also GSC 04766–01466.
* HD 36780 is a K5III giant as tabulated in SIMBAD. It has a (variable) radial velocity inconsistent with association membership ($V_r = +91$ km s^{-1}, Griffin 1972; $V_r = +73.2$ km s^{-1}, Bois et al. 1988). The star has an *Hipparcos* distance of 260±50 pc.
* HD 290675 has a discordant radial velocity (Gieseking 1983). It is also BD–01 967.
* HD 37491 is likely associated to the mid-infrared source IRAS 05363–0111.
* HD 290749 is a B8V star according to Nesterov et al. (1995). With proper motion of 13.8 mas a^{-1} and colour $V_T - K_s \sim 1.0$ mag, it is likely a late A- or an early F-type star in the foreground. See also the note for TYC 4766 2150 1 (Table A.5).
* HD 36882 is at $d = 220 \pm 50$ pc (Perryman et al. 1997). Because of a transcription error, SIMBAD tabulates HD 36882 as one of the early-type stars associated to the H II region Sh 2–264, close to λ Ori, in Sharpless (1959); the actual early-type star is ϕ^{01} Ori (HD 36822, B0III).
* HD 290667 and StHA 46 were catalogued by Stephenson (1986) as Hα emission stars. However, Downes & Keyes (1988) and Maheswar et al. (2003) failed to detect the (sporadic?) emission. StHA 46 is at 18 arcsec to the southwest of the early-A star AG–00 669.
* HD 290647 falls in the tiny overlapping region between the Alnilam and Mintaka fields. It is also BD–00 1001.
* TYC 4767 2257 1 has a colour $V_T - K_s > 4.5$ mag, typical of intermediate M stars. Its proper motion is, however, very low ($\mu \lesssim 1$ mas s^{-1}).

Notes to Table A.8:

* HD 290647 falls in the tiny overlapping region between Alnilam and Mintaka fields. It is also BD–00 1001.
* TYC 4766 516 1 has a very red colour of $V_T - K_s = 5.44 \pm 0.09$ mag and no *IRAS* excess. It could be a mid-M-type giant or subgiant in back-/foreground.
* HD 290576 is also BPM 71736.
* TYC 4766 2124 1 is also GSC 04766–02124.
* HD 290568 is also BD–00 987 and IRAS 05303–0009.
* HD 36117 is a nearby ($d = 170 \pm 30$ pc; Perryman et al. 1997), peculiar, A-type star (Gray & Corbally 1993) with X-ray emission
* HD 36139 is a nearby ($d = 124 \pm 13$ pc; Perryman et al. 1997), high rotation-velocity, radial-velocity variable (Morrell & Levato 1991), A-type star with no known companion.
* HD 36840 is at an *Hipparcos* distance of $d = 380 \pm 120$ pc, which is probably incorrect, given the spectral type of the star (G5V).
* HD 36558 has a discordant radial velocity of $V_r = +42.1 \pm 4.8$ km s^{-1} (Nordström et al. 2004).
* HD 36443 (LHS 5107, G 99–16; Roman 1995) is a well-known, solar-like, high-velocity star at only $d = 38.2 \pm 1.9$ pc and with radial velocity $V_r \approx -9.1$ km s^{-1} (Wilson 1953).
* HD 290486 AB is a visual binary star with $\rho \approx 1.7$ arcsec, $\theta \approx 304$ deg (Dommanget & Nys 1994).

Notes to Table A.9:

* E Ori 2–1328 is a young M4.5-type very low-mass star with lithium in absorption (pEW(Li I) = +0.40±0.05 Å) and Balmer lines in faint (chromospheric) emission (pEW(Hα) = –8.2±0.5 Å). The sodium line in the red optical is weak in comparison with field dwarfs of the same spectral type (pEW(Na I) = +3.4±0.5 Å; Béjar et al. 2003a).
* E Ori 2–1868 is a young M6.0-type very low-mass star with faint alkali lines (pEW(Na I) < 3 Å; Béjar et al. 2003b). The *i*-band magnitude has been taken from the SuperCOSMOS Science Archive (Hambly et al. 2001).
* E Ori 1–388 is a young M6.0-type very low-mass star with faint alkali lines (pEW(Na I) = +3.9±0.5 Å; Béjar et al. 2003a) and Balmer lines in faint (chromospheric) emission (pEW(Hα) = –6.5±2.0 Å). It is embedded in the [OS98] 40B remnant molecular cloud.
* E Ori 1–1644 is a young M5.0-type very low-mass star with Balmer lines in faint (chromospheric) emission (pEW(Hα) = –7.6±1.0 Å; Béjar et al. 2003a).
* E Ori 2–878 is a young M5.5-type very low-mass star with faint alkali lines (pEW(Na I) < 3 Å; Béjar et al. 2003b).
* E Ori 2–705 is a young M5.0-type very low-mass star with faint alkali lines (pEW(Na I) < 4 Å; Béjar et al. 2003b). The *i*-band magnitude is from SuperCOSMOS.
* E Ori 2–603 is a young M5.5-type very low-mass star with faint alkali lines (pEW(Na I) < 3 Å; Béjar et al. 2003b).
* Kiso A–0904 41 and Kiso A–0904 42 form a binary system with $\rho \approx 11.0$ arcsec, $\theta \approx 277$ deg. The *i*-band magnitudes are from SuperCOSMOS. They hay been identified in *XMM-Newton* observations (Caballero et al., in prep).

* V469 Ori is proably associated to the [OS98] 29J, [OS98] 29H and [OS98] 29K remnant molecular clouds.
* Kiso A–0904 76 is a K6-type variable (ΔV = 0.33 mag) star with pEW(Hα) = –20.2 Å and pEW(Li I) = +0.5 Å (Briceño et al. 2005). It has a visual companion at $\rho \approx 5.3$ arcsec, $\theta \approx 321$ deg ($\Delta H = 2.91 \pm 0.05$ mag).
* Haro 5–80 is a variable, emission-line star (Haro & Moreno 1953; Fedorovich 1960; Wiramihardja et al. 1989). It has a very nearby ($\rho \sim 3$ arcsec, $\theta \sim 15$ deg) companion or small jet (possibly associated to an unknown Herbig-Haro object)
* 2E 1398 is an X-ray source tabulated in at least six catalogs from *Einstein*, *ROSAT* and *XMM-Newton* data (Harris et al. 1994; McDowell 1994; Moran et al. 1996; Voges et al. 1999; *ROSAT* Consortium 2000; *XMM-Newton* Survey Science Centre Consortium 2007). It is located at 4.5 arcmin to the west of Alnilam. The *i*-band magnitude is from SuperCOSMOS.
* Kiso A–0904 37 has rather blue $I - J$ and $J - K_s$ colours. Besides, it has a faint, red, visual companion at $\rho \approx 5.6$ arcsec, $\theta \approx 144$ deg, unresolved by Wiramihardja et al. (1989). It is likely that the actual emission-line star or brown dwarf (and the only truly young objects) is the visual companion (J2000 coordinates: 05 35 32.38 –01 12 08.2).
* StHA 47 is a mid-K-type T Tauri star according to Downes & Keyes (1988). It is the fourth strongest X-ray emitter at less than 20 arcmin to Alnilam, from *XMM-Newton* observations.
* CVSO 162 is an M1-type variable (ΔV = 0.23 mag) star with pEW(Hα) = –3.9 Å and pEW(Li I) = +0.5 Å (Briceño et al. 2005).
* Annizam 363062 is a visual binary with $\rho \approx 6.3$ arcsec, $\theta \approx 243$ deg ($\Delta H = 2.41 \pm 0.06$ mag). Five X-ray events in the surrounding area were tabulated in the catalogue of *ROSAT* HRI Pointed Observations (ROSAT Team 2000). It might also be the *Einstein* source 2E B0534–0111. It is not known which component is the actual X-ray emitter.
* Haro 5–67 is a G:-type, photometrically variable, T Tauri star with strong Balmer emission detected by several Hα objective-prism surveys (Haro & Moreno 1953; Sanduleak 1971; Stephenson 1986) and with *IRAS* flux excess (Weaver & Jones 1992). It has been spectroscopically followed-up by Herbig & Kaneswara Rao (1972) and Downes & Keyes (1988).
* Kiso A–0904 50 has rather blue $I - J$ and $J - K_s$ colours. It might be a variable young star or an active object in the fore-/background.
* Kiso A–0904 61 could also be the Hα emitter Haro 5–77 (suspected variable NSV 2465; Kukarkin et al. 1981).
* V583 Ori (Haro 5–74) is a variable, emission-line star (Haro & Moreno 1953; Fedorovich 1960; Wiramihardja et al. 1989). It is possibly the X-ray source 2E 1423 and might form a Lindroos system together with the B9V star HD 37187 (Table A.3).
* HD 36779 B is a K5IV with Li I in absorption in a likely Lindroos system with the B-type star HD 36779 (Table A.3 – Lindroos 1985; Pallavicini, Pasquini & Randich 1992; Martín, Magazzù & Rebolo 1992). See, however, a brief discussion on the position in the H-R diagram and the radial velocity of HD 36779 B in Gerbaldi, Faraggiana & Balin (2001).
* 2E 1357 is a K3-type star with pEW(Hα) = –2.75 Å and pEW(Li I) = +0.470±0.008 Å (Alcalá et al. 1996, 2000).
* Kiso A–0904 28 is also the X-ray source 2E 1340 (McDowell 1994). The star was detected by Kraemer et al. (2003) at 8.3 μm, which suggests a possible flux excess in the mid-infrared.
* Kiso A–0904 30 was previously identified with a fainter, much bluer source 24 arcsec to the east.
* 2E 1449 is a K4-type star with pEW(Hα) = +0.35 Å and pEW(Li I) = +0.410±0.010 Å (Alcalá et al. 1996, 2000).
* Kiso A–0904 60 is a K6-type variable (ΔV = 0.51 mag) star with pEW(Hα) = –61.9 Å and pEW(Li I) = +0.3 Å (Briceño et al. 2005). It could be a tight binary ($\rho \lesssim 1.0$ arcsec) based on the 2MASS photometry quality flags.
* Kiso A–0904 65 is a variable (ΔV = 0.87 mag) star with a very strong Balmer emission, pEW(Hα) = –400 Å, and lithium in absorption, pEW(Li I) = +0.3 Å (Briceño et al. 2005).
* CVSO 124 is an M3-type variable (ΔV = 0.19 mag) star with pEW(Hα) = –17.3 Å and pEW(Li I) = +0.4 Å (Briceño et al. 2005). It also falls in the Mintaka field.
* Kiso A–0904 34 is also [SE2005] 104. It does not show significant periodic variability.
* Kiso A–0904 33 has a faint, red, visual companion at $\rho \approx 4.9$ arcsec, $\theta \approx 14$ deg, with near-infrared magnitudes $J = 12.55 \pm 0.03$ mag and $K_s = 11.55 \pm 0.03$ mag.
* PU Ori is a pre-main sequence star with Hα in strong two-lobe emission (Haro & Moreno 1953; Herbig & Kameswara Rao 1972; Cohen & Kuhi 1979; Wiramihardja et al. 1989), photometric variability (Fedorovich 1960; Briceño et al. 2005), mid-infrared flux excess at the *IRAS* passbands (Weintraub 1990; Weaver & Jones 1992) and forbidden emission lines ([O I] λ 6300.3 Å; Hirth, Mundt & Solf 1997). It has an extraordinary red colour of $J - K_s = 1.77 \pm 0.03$ mag.
* StHA 48 is a K4-type T Tauri star according to Maheswar et al. (2003).

Notes to Table A.10:

* CVSO 124 is an M3-type variable (ΔV = 0.19 mag) star with pEW(Hα) = –17.3 Å and pEW(Li I) = +0.4 Å (Briceño et al. 2005). It also falls in the Alnilam field.
* Kiso A–0903 221 has a faint red visual companion at $\rho \approx 6.4$ arcsec, $\theta \approx 49$ deg ($\Delta H = 5.1 \pm 0.2$ mag).
* Kiso A–0904 4 is a K7-type variable (ΔV = 0.66 mag) star with pEW(Hα) = –34.2 Å and pEW(Li I) = +0.3 Å (Briceño et al. 2005). It is also Kiso A–0903 228.
* IRAS 05307–0038 (also known as YSO CB031YC1 and YSO–C CB031YC1–I) is a bright ($K_s = 8.63 \pm 0.02$ mag) T Tauri star embedded in the IC 434 Bok globule/reflection nebula (Dreyer 1895; Hubble 1922; Cederblad 1946; Dorschner & Gürtler 1963; Magakian 2003). The star has an extended and fuzzy nebulosity in the *J* band and was classified as a Class II object because of a flux excess at 25 μm (Yun & Clemens 1994, 1995). Yun et al. (1997) mea-

sured Hα and Hβ in emission and Li I in absorption, and characterised its SED from the *B* band to the mid-infrared. Yun et al. (1996) discovered two nearby radio sources that could be associated to the star (there are other two additional radio sources in the surrounding area, found by Condon et al. 1998). Gómez et al. (2006), in a very sensitive survey using NASA 70 m antenna at Robledo de Chavela (Spain), failed to detect water maser emission from these sources. See Fig. A.1.

* Kiso A–0904 22 has a nearby visual companion identified as a photometric young star candidate, Mantaqah 2385113 (Table A.17).
* Kiso A–0904 13 is also Kiso A–0903 245.
* Mantaqah 487126 displayed 14 X-ray events during *ROSAT* observations (White et al. 2000 –8 events–; ROSAT 2000 –6 events–).
* 1AXG J053127–0021 appears in numerous X-ray catalogs from *ROSAT* and *ASCA* data. The *i*-band magnitude is from SuperCOSMOS.
* Mantaqah 148186 is an X-ray source in several catalogs (e.g. Voges et al. 1999).
* Mantaqah 400105 is an X-ray source in several catalogs (e.g. White et al. 2000).
* Mantaqah 320042 displayed 12 X-ray events during *ROSAT* observations (White et al. 2000 –5 events–; ROSAT 2000 –7 events–).
* Kiso A–0904 18 has a $J - K_s$ colour redder than 2.0 mag.

Notes to Table A.11:

* E Ori 2–1982 has no DENIS or SuperCOSMOS counterpart. The *i*-band magnitude (actually *I*) is from Béjar et al. (2003b).
* [OSP2002] OriI–2N 4 has a tiny (chromospheric?) Balmer emission (pEW(Hα) $\approx$ –2.6 Å; Ogura, Sugitani & Pickles 2002).
* BD–01 969B ("Alnilam B", ϵ Ori B; V = 10.5 mag) is located at $\rho \approx 179.0$ arcsec, $\theta \approx 58$ deg, to Alnilam. (this value coincides with the original measurements by Burnham in 1879 – Burnham 1906). No spectral type has been tabulated or measured.
* AG–00 669 forms a visual double with the foreground solar-like star StHA 46 (Jeffers, van den Bos & Greeby 1963). See Table A.7.
* BD–00 983B ("Mintaka B", δ Ori B; V = 14.0 mag) is located at $\rho \approx 33.0$ arcsec, $\theta \approx 229$ deg, to Mintaka AE–D (this value coincides with the original measurements by Burnham in 1878 – Burnham 1906). No spectral type has been tabulated or measured. The star shows no evidence of any significan X-ray emission in deep observations with the *Chandra* Space Telescope (Miller et al. 2002).
* BD–00 993B ("HD 36726 BC"; V = 13.7 mag) is located at $\rho \sim 19.8$ arcsec, $\theta \sim 213$ deg ($\Delta H = 1.30 \pm 0.05$ mag) to the A0Vm-type star HD 36726 (Table A.4). BD–00 993B is, in its turn, a close binary with $\rho \sim 0.8$ arcsec, $\theta \sim 64$ deg (Dommanget & Nys 1994).

Notes to Table A.12:

* X Ori is the reddest object in the studied area. It is an M8–9-type Mira Cet variable found by Wolf (1904) with P = 424.15±1.77 d (Templeton, Mattei & Willson 2005) and silicate dust emission (Sloan & Price 1998; Speck et al. 2000). Although X Ori is even brighter in the near- and mid-infrared than Alnilam and Mintaka ($K_s \sim 0.9$ mag), its extremely red $V-K_s$ colour, of more than 10 mag, prevented its detection in the Tycho-2 catalogue.
* Ruber 1 has a proper motion of –12±5, –49±2 mas s^{-1}, measured by us using POSSI, UKST blue, red and infrared, DENIS and 2MASS (the six epochs cover 47 years; see method details in Caballero 2007b). Ruber 1 probably is a late M dwarf in the foreground.
* G 99–18 is a high proper-motion star, with μ = 280 mas a^{-1}. It falls in the tiny overlapping region between the Alnilam and Mintaka fields.
* Ruber 2 has very peculiar colours: the near-infrared colours are very red (e.g. $J - K_s = 1.33 \pm 0.04$ mag), typical of early and intermediate L dwarfs, very late M giants or carbon stars. Its $I - J$ colour is red enough, as well, to be selected as an association member candidate. The optical colours are, however, contradictory and variable. The SuperCOSMOS Science Archive tabulates photographic magnitudes B(1988.0) = 17.643 mag, R_1(1951.9) = 19.148 mag and R_2(1989.0) = 15.713 mag. Photographic *R* bands are separated by 37.1 years and probably reflect intrinsic (high-amplitude, long-time-scale) photometric variability of the object. SuperCOSMOS and USNO-B1 tabulate appreciable proper motions for Ruber 2. However, after a careful astrometric study using the original plate digitisations, DENIS and 2MASS, we conclude that the proper motion of the object is null within uncertainties of 10 mas s^{-1}. Ruber 2 is at a separation of only ∼9 arcmin to the core of the unusual Berkeley 20 cluster (see Section 1). Therefore, Ruber 2 probably is a pulsating late M giant of that cluster, at an heliocentric distance of 8.4 kpc.

Notes to Table A.14:

* 2MASS J05345451–0143256 has a 2MASS double detection, with quality flags AUU and UEA. It is not in the NASA/IPAC Extragalactic Database (NED). It could be an unresolved stellar binary instead of a galaxy.
* 2MASX J05332498–0106242 is also the NED object LCSB S0895N (Monnier Ragaigne et al. 2003).
* 2MASX J05365804–0042413 is the infrared source IRAS 05344–0044, very close to the young early-type star HD 37235.
* 2MASX J05364723–0039144 is also NED object LCSB S0899N (Monnier Ragaigne et al. 2003).
* 2MASS J05364746–0039110 is located at 4.3 arcsec to the centre of 2MASX J05364723–0039144 (see just above), in the plane of the galaxy. This source is probably an artifact.

Notes to Table A.15:

Fig. A.1. False-colour composite image, 7.5×7.5 arcmin^2 wide, of the IC 434 Bok globule, centred on IRAS 05307–0038 (labelled as YSO CB031YC1). Blue, green and red are for photographic B_J, R_F and I_N, respectively. North is up, east is left. The six reddened sources in the IC 434 Bok globule (Table A.13) are indicated with (red) circles. *[NOTE to the reader: this image is only available at A&A].*

* 2MASX J05322266–0000555 is also the NED object LEDA 147610 (Klemola, Jones & Hanson 1987; Paturel et al. 1989).
* 2MASS J05341337–0044087 is PMN J0534–0044, a powerful radio source discovered in many surveys (e.g. Becker, White & Edwards 1991; Griffith et al. 1995; Douglas et al. 1996; Condon et al. 1998). Its optical/near infrared counterpart is faint ($i = 16.82 \pm 0.11$ mag) and relatively blue ($i - K_s = 1.2 \pm 0.2$ mag).

Notes to Table A.16:

* Annizam 2473146 is at ∼4 arcsec to the east of an extended source with a galactic appearance.
* Annizam 2464138 could be reddened by the nearby Ori I–2 Bok globule and be a background star.
* Annizam 1840146 was subject of a dedicated astrometric study using public data (plate digitisations, DENIS and 2MASS). There seems to be an artifact in the POSSI Schmidt plate of 1951 that causes a false proper motion of more than 100 mas s^{-1} in the astrometric catalogs USNO-B1 and SuperCOSMOS Science Archive. The very red object has, however, null proper motion within the uncertainty of 10 mas s^{-1} using seven astrometric epochs between 1987 and 2000 (see Caballero 2007b for details of the astrometric analysis).
* Annizam 1106127 has 2MASS photometry quality flags "EEE"; it might indicate that it is an unresolved binary with $\rho \sim 1$–2 arcsec.
* Annizam 1415101 is surrounded by a galaxy arm-like structure. It could be the point-like core of a background galaxy.
* Annizam 2042095 has a bluer, fainter, visual companion at about 3 arcsec to the northeast.
* Annizam 2611268 is at ∼8 arcsec to the south of a probable foreground star, about ∼3.3 mag brighter in the *i* band.
* Annizam 123132 is in the glare of Alnilam ($\rho \sim 2.0$ arcmin, $\theta \sim 130$ deg).
* Annizam 1751268 is probably the *Einstein* X-ray source 2E 1458.
* Annizam 1748267 is at only $\rho \sim 8.4$ arcsec, $\theta \sim 340$ deg to Annizam 1751268 (see just above).
* V993 Ori is a long-time-known variable star discovered by Luyten (1932). It has a very red colour $J - K_s = 1.26 \pm 0.04$ mag for its brightness ($H = 9.82 \pm 0.02$ mag).
* Annizam 798196 could be associated to the X-ray source 1WGA J0536.4–0059, found in several *ROSAT* HRI and PSPC catalogues (Moran et al. 1996; *ROSAT* Consortium 2000; Flesch & Hardcastle 2004). It might be a background source associated to the (extragalactic?) radio source NVSS 053627–005937 in the 1.4 GHz NRAO VLA Sky Survey (Condon et al. 1998), located at $\rho \sim 18$ arcsec, $\theta \sim 200$ deg.
* [SE2005] 120 (Mantaqah 1357158) is a photometric candidate member of the "ϵ Orionis cluster" firstly identified by Scholz & Eislöffel (2005). It is the most variable star of the five objects with high-amplitude ($A_I = 0.952$ mag), irregular variations. It also have a significant periodic variability of $P = 82 \pm 3$ h, with superimposed short-term fluctuations. Scholz & Eislöffel (2005) classified it as very low-mass analogue of classical T Tau stars affected by intrinsic reddening. For explaining the large variations, they proposed two possible scenarios involving a eclipsing "hot Jupiter" in close orbit and "hot spots formed by matter flow from an accretion disc onto the central object". It might be, however, a typical eclipsing binary in the fore-/background.
* [SE2005] 126 (Mantaqah 1582164) is other photometric candidate member of the "ϵ Orionis cluster" with significant periodic variability in the work by Scholz & Eislöffel (2005). In this case, the object has a low-amplitude variability ($A_I = 0.016$ mag) with a very short period ($P = 4.06 \pm 0.05$ mag). The values are consistent with pulsations induced by deuterium-burning in young brown dwarfs (Caballero et al. 2004; Palla & Baraffe 2005).
* [SE2005] 71 (Annizam 2446149) is a non-variable photometric candidate member of the "ϵ Orionis cluster" (Scholz & Eislöffel 2005). It is located at $\rho \sim 14$ arcsec, $\theta \sim 270$ deg, to V472 Ori.

Notes to Table A.17:

* Mantaqah 2041159 is surrounded by galaxy arm-like structures. It could be the point-like core of a background galaxy.
* [SE2005] 44 (Mantaqah 2627125) is a photometric candidate member of the "ϵ Orionis cluster" with significant periodic variability ($A_I = 0.027$ mag, $P = 31.0 \pm 1.8$ h; Scholz & Eislöffel 2005).
* Mantaqah 2216132 has a fainter redder visual companion at $\rho \sim 5.6$ arcsec, $\theta \sim 210$ deg.
* Mantaqah 2385113 is at $\rho \approx 6.87$ arcsec, $\theta \approx 352$ deg to Kiso A–0904 22, together with it could form a ∼2700 AU-wide low-mass binary.
* Mantaqah 941101 has a brighter bluer visual companion at $\rho \sim 4.5$ arcsec, $\theta \sim 230$ deg.
* Mantaqah 1926266 is at $\rho \approx 15.1$ arcmin, $\theta \approx 224$ deg, to Kiso A–0903 183.
* Mantaqah 161138 is in the glare of Mintaka ($\rho \sim 2.7$ arcmin, $\theta \sim 140$ deg).
* Mantaqah 104142 is in the glare of Mintaka ($\rho \sim 1.7$ arcmin, $\theta \sim 140$ deg).
* Mantaqah 67144 is in the glare of Mintaka ($\rho \sim 1.1$ arcmin, $\theta \sim 145$ deg). Its *J*-band magnitude is strongly affected.
* Mantaqah 1982094 has a blue visual companion at $\rho \sim 6.6$ arcsec, $\theta \sim 60$ deg.

Notes to Table A.18:

* Albus 1 has by far the bluest colour among all the investigated objects ($V_T - K_s = -0.95 \pm 0.14$ mag) and a measurable proper motion $\mu = 19$ mas a^{-1}. Although Albus 1 was detected from the data presented in this work, it was followed-up with additional photometric data and discussed in detail in Caballero & Solano (2007). Very recently, Vennes, Kawka & Allyn Smith (2007) obtained a series of optical spectra, showing that it is a peculiar, bright, helium-rich B3 subdwarf ($n_{He} = 0.6 \pm 0.1$).
* TYC 5360 681 1, the stellar counterpart of an *IRAS* source, has an appreciable proper motion tabulated by Tycho-2: ($\mu_\alpha \cos\delta$, μ_δ) = (–4±2, +16±3) mas a^{-1}. Assuming an heliocentric distance of $d \gtrsim 1$ kpc, typical of a giant, it would have a very large tangential velocity of $V_t \gtrsim 80$ km s^{-1}.
* BD–13 1293 is the reddest *IRAS* source in this list. The star, investigated here for the first time, displays very strong wide absorption bands, especially at $\sim 1\,\mu$m (between the *i* and *J* bands). It is a photometric variable: from the Hipparcos catalogue, its V_T magnitude varies between V_T = 9.41 and 10.00 mag (15 and 85 % percentiles, respectively).

Table A.1. Photometry of correlated Tycho-2/2MASS sources in the Alnilam field.

Name	TYC	B_T (mag)	V_T (mag)	$J \pm \delta J$ (mag)	$H \pm \delta H$ (mag)	$K_s \pm \delta K_s$ (mag)
HD 290690	4770 1664 1	11.423	10.847	9.771 0.023	9.513 0.033	9.445 0.021
...	4770 1072 1	11.747	10.271	8.056 0.018	7.479 0.034	7.339 0.018
HD 36980 AB	4770 1635 1	9.044	9.007	8.797 0.024	8.898 0.042	8.812 0.019
RX J0535.6–0152 AB	4770 1685 1	11.660	12.113	10.541 0.022	10.223 0.028	10.167 0.021
...	4766 2029 1	12.161	10.987	9.678 0.023	9.240 0.027	9.127 0.019
HD 290691	4766 1575 1	10.505	10.244	9.926 0.024	9.817 0.030	9.821 0.021
...	4767 3 1	12.214	10.230	7.132 0.023	6.381 0.031	6.111 0.020
HD 290686	4766 1897 1	11.290	11.048	10.372 0.019	10.343 0.031	10.221 0.023
HD 37038 AB	4766 1653 1	8.542	8.157	7.348 0.019	7.183 0.023	7.150 0.018
...	4766 1605 1	12.112	11.704	10.628 0.022	10.331 0.028	10.259 0.023
...	4767 935 1	12.069	11.737	10.475 0.028	10.213 0.024	10.138 0.021
HD 290684	4766 1617 1	9.723	9.651	9.382 0.023	9.366 0.032	9.359 0.018
...	4766 1891 1	12.273	11.700	10.388 0.021	10.051 0.030	9.959 0.019
HD 37285 AB	4766 1893 1	9.082	8.983	8.839 0.027	8.873 0.030	8.810 0.024
HD 37113	4766 1493 1	8.665	8.652	8.525 0.020	8.484 0.031	8.518 0.020
HD 290685	4766 1923 1	10.890	9.584	7.396 0.023	6.833 0.024	6.728 0.024
HD 37389	4767 1118 1	8.251	8.325	8.477 0.029	8.551 0.033	8.568 0.024
HD 36863	4766 119 1	8.585	8.280	7.715 0.019	7.684 0.027	7.635 0.018
...	4766 1699 1	11.411	10.983	10.074 0.019	9.965 0.028	9.919 0.020
HD 290683	4766 1993 1	11.942	11.593	10.424 0.023	10.173 0.032	10.128 0.020
IRAS 05354–0142	4767 829 1	12.531	11.115	5.152 0.018	4.308 0.076	3.972 0.320
...	4766 121 1	12.328	11.461	10.259 0.024	9.971 0.032	9.900 0.021
HD 290682	4766 2001 1	10.310	9.764	8.675 0.030	8.474 0.049	8.388 0.023
...	4766 1823 1	11.795	11.489	10.493 0.019	10.317 0.030	10.254 0.022
HD 37272	4766 1809 1	7.752	7.844	8.041 0.023	8.101 0.034	8.028 0.027
...	4766 400 1	12.268	11.598	11.037 0.022	10.867 0.032	10.767 0.023
...	4766 1643 1	11.786	10.818	8.549 0.023	7.918 0.029	7.804 0.021
HD 37149	4766 1903 1	7.898	7.989	8.162 0.027	8.299 0.038	8.296 0.026
HD 290770	4766 2027 1	9.198	9.238	8.570 0.018	7.902 0.034	7.078 0.023
HD 290681	4766 1705 1	10.269	9.836	8.815 0.024	8.581 0.040	8.521 0.022
...	4766 2063 1	12.665	11.416	8.968 0.021	8.266 0.020	8.116 0.034
HD 37344	4767 1011 1	8.703	8.712	8.809 0.048	8.783 0.038	8.805 0.019
HD 290679	4766 1466 1	11.566	10.258	7.828 0.023	7.264 0.053	7.115 0.018
HD 290680	4766 1556 1	11.097	10.616	9.946 0.022	9.825 0.030	9.720 0.022
...	4766 260 1	12.183	11.668	10.976 0.024	10.809 0.031	10.780 0.021
...	4766 265 1	12.866	11.298	9.345 0.022	8.729 0.030	8.572 0.020
HD 290602	4766 211 1	10.547	10.303	10.049 0.022	10.031 0.030	9.998 0.019
...	4766 1599 1	11.176	10.760	9.698 0.022	9.458 0.033	9.439 0.021
HD 290678	4766 1490 1	10.801	10.706	10.161 0.022	10.118 0.032	10.039 0.019
...	4767 498 1	12.154	11.308	8.970 0.046	8.368 0.055	8.184 0.021
[NYS99] A–06	4766 108 1	11.943	11.565	10.527 0.022	10.098 0.030	9.980 0.023
HD 290677	4766 1535 1	10.271	10.087	9.574 0.022	9.469 0.030	9.420 0.019
HD 36780	4766 1915 1	7.930	6.080	3.445 0.222	2.623 0.198	2.401 0.280
...	4766 144 1	11.993	11.660	10.407 0.022	10.120 0.031	10.068 0.021
...	4766 1833 1	12.035	12.084	10.761 0.024	10.562 0.031	10.486 0.020
HD 290674	4766 1797 1	9.860	9.745	9.558 0.021	9.519 0.033	9.482 0.023
...	4766 2326 1	11.736	10.433	8.333 0.023	7.785 0.055	7.671 0.021
HD 37321 AB	4767 271 1+2	7.0558	7.2085	7.281 0.039	7.252 0.040	7.243 0.020
HD 290673	4766 1689 1	10.108	9.822	9.054 0.023	8.920 0.030	8.876 0.021
HD 36955	4766 330 1	9.586	9.451	9.141 0.021	9.077 0.031	9.071 0.021
HD 290675	4766 2413 1	9.693	9.192	8.193 0.021	8.003 0.029	7.942 0.023
HD 290767	4767 851 1	11.527	10.156	7.853 0.029	7.295 0.036	7.161 0.029
HD 290766	4767 828 1	10.905	10.642	9.990 0.027	9.923 0.026	9.847 0.022

Table A.1. Photometry of correlated Tycho-2/2MASS sources in the Alnilam field (cont.).

Name	TYC	B_T (mag)	V_T (mag)	$J \pm \delta J$ (mag)	$H \pm \delta H$ (mag)	$K_s \pm \delta K_s$ (mag)
...	4767 1130 1	12.876	12.095	10.913 0.064	10.648 0.026	10.555 0.026
HD 290676	4766 2409 1	10.411	9.859	8.922 0.018	8.752 0.028	8.696 0.023
HD 290672	4766 2375 1	10.464	10.171	9.784 0.022	9.702 0.031	9.654 0.019
HD 290765	4767 671 1	9.074	9.017	9.005 0.027	8.948 0.042	8.955 0.023
V1247 Ori	4767 953 1	10.177	9.852	8.878 0.032	8.203 0.053	7.408 0.029
HD 37172	4766 2451 1	8.457	8.329	7.993 0.019	7.988 0.046	7.926 0.020
...	4766 2371 1	11.238	11.038	10.320 0.024	10.243 0.031	10.164 0.019
HD 37506	4767 473 1	9.038	8.553	7.648 0.029	7.490 0.038	7.399 0.026
...	4766 2344 1	11.969	11.278	10.206 0.026	9.934 0.022	9.881 0.021
...	4766 2221 1	12.198	12.270	11.150 0.026	11.022 0.022	10.941 0.019
...	4766 724 1	11.136	10.692	9.536 0.023	9.235 0.026	9.194 0.021
Alnilam A	4766 2450 1	1.553	1.692	2.191 0.324	2.408 0.180	2.273 0.282
...	4767 487 1	12.828	11.639	9.568 0.027	9.025 0.024	8.909 0.024
HD 37491	4767 897 1	8.972	7.661	5.727 0.029	5.205 0.042	5.020 0.017
HD 37397	4767 871 1	6.637	6.802	7.163 0.029	7.225 0.057	7.259 0.021
VV Ori	4766 2449 1	5.144	5.340	5.754 0.023	5.815 0.038	5.862 0.027
HD 290763	4767 675 1	11.258	10.586	9.508 0.027	9.235 0.027	9.169 0.027
...	4766 477 1	11.097	11.162	10.498 0.026	10.201 0.023	10.131 0.021
HD 36684 AB	4766 507 1	8.621	8.636	8.548 0.021	8.602 0.059	8.546 0.019
HD 37443	4767 433 1	8.410	7.827	6.843 0.029	6.665 0.029	6.555 0.024
...	4766 1854 1	11.018	10.757	9.952 0.024	9.801 0.024	9.791 0.021
HD 290670	4766 2362 1	10.442	9.820	8.806 0.023	8.550 0.047	8.525 0.021
HD 290750	4767 1235 1	10.118	9.805	9.477 0.026	9.430 0.024	9.379 0.024
...	4766 2150 1	11.695	11.322	10.292 0.027	10.086 0.026	10.016 0.023
HD 36779 AC	4766 559 1	6.013	6.196	6.575 0.019	6.630 0.036	6.701 0.018
HD 36996	4766 2250 1	9.567	9.508	9.476 0.023	9.486 0.033	9.462 0.021
HD 37187	4766 2448 1	8.112	8.128	8.054 0.027	8.054 0.034	8.016 0.016
HD 290671	4766 2458 1	8.906	8.955	8.871 0.021	8.824 0.031	8.757 0.018
HD 37076	4766 2447 1	7.943	8.022	8.156 0.023	8.209 0.034	8.230 0.024
HD 290753	4767 1217 1	11.234	10.890	10.021 0.026	9.798 0.024	9.721 0.019
...	4766 1756 1	12.312	11.929	11.001 0.030	10.691 0.029	10.630 0.030
HD 36882	4766 802 1	8.392	7.108	5.252 0.228	4.482 0.036	4.243 0.020
...	4767 760 1	12.244	11.114	9.221 0.032	8.756 0.046	8.603 0.028
HD 290751	4767 511 1	11.816	11.371	10.532 0.026	10.355 0.023	10.266 0.021
HD 290748	4766 1874 1	10.331	10.233	9.918 0.027	9.840 0.026	9.850 0.021
HD 37075	4766 2165 1	9.382	9.347	9.278 0.023	9.267 0.030	9.279 0.020
HD 37302	4767 556 1	9.052	9.071	9.130 0.029	9.153 0.023	9.122 0.019
HD 290666	4766 750 1	9.813	9.738	9.613 0.024	9.592 0.032	9.578 0.023
...	4767 1008 1	12.234	11.076	8.557 0.023	7.881 0.055	7.745 0.031
HD 290664	4766 1906 1	10.639	10.016	8.892 0.024	8.588 0.034	8.589 0.028
HD 290665	4766 656 1	9.523	9.425	9.175 0.023	9.196 0.032	9.149 0.023
HD 36915	4766 608 1	7.960	7.985	7.996 0.058	7.986 0.024	7.981 0.024
HD 290752	4767 744 1	11.485	10.779	9.596 0.028	9.282 0.024	9.178 0.023
HD 290663	4766 2204 1	10.452	10.147	9.591 0.024	9.428 0.032	9.369 0.019
HD 37112	4766 2177 1	7.905	7.970	8.110 0.037	8.145 0.055	8.164 0.024
V1379 Ori	4767 649 1	7.458	7.573	7.906 0.030	7.953 0.042	8.003 0.040
HD 290662	4766 2211 1	9.920	9.817	9.389 0.023	9.261 0.030	9.206 0.020
HD 36825	4766 525 1	8.684	8.616	8.581 0.021	8.615 0.049	8.608 0.021
HD 36954 AB	4766 2058 1	6.817	6.955	7.116 0.027	7.150 0.026	7.137 0.021
...	4767 715 1	12.740	11.473	9.252 0.030	8.602 0.038	8.475 0.023
HD 37235	4766 2242 1	8.029	8.131	8.305 0.024	8.387 0.038	8.382 0.019
HD 37284	4766 1411 1	9.089	8.931	8.757 0.026	8.717 0.042	8.687 0.021
HD 290746	4767 607 1	11.775	11.609	10.643 0.023	10.449 0.023	10.397 0.019

Table A.1. Photometry of correlated Tycho-2/2MASS sources in the Alnilam field (cont.).

Name	TYC	B_T (mag)	V_T (mag)	$J \pm \delta J$ (mag)	$H \pm \delta H$ (mag)	$K_s \pm \delta K_s$ (mag)
...	4766 2257 1	13.837	11.545	7.908 0.027	7.068 0.046	6.743 0.027
HD 290648	4766 1170 1	9.726	9.604	9.510 0.022	9.474 0.025	9.435 0.023
HD 290660	4766 1836 1	10.825	10.578	10.109 0.024	10.063 0.040	9.940 0.024
HD 290745	4767 371 1	10.807	10.525	9.933 0.026	9.773 0.023	9.743 0.023
HD 290650	4766 1968 1	10.415	10.247	9.757 0.022	9.668 0.032	9.593 0.018

Table A.2. Photometry of correlated Tycho-2/2MASS sources in the Mintaka field.

Name	TYC	B_T (mag)	V_T (mag)	$J \pm \delta J$ (mag)	$H \pm \delta H$ (mag)	$K_s \pm \delta K_s$ (mag)
...	4766 590 1	11.055	10.325	9.108 0.020	8.841 0.059	8.722 0.019
HD 290585	4766 602 1	11.329	11.012	10.355 0.029	10.241 0.035	10.196 0.029
...	4766 644 1	12.670	11.104	8.727 0.037	8.035 0.034	7.870 0.024
...	4766 687 1	13.049	11.051	8.192 0.020	7.439 0.059	7.183 0.027
...	4766 733 1	12.498	12.726	10.954 0.023	10.651 0.026	10.575 0.023
...	4766 744 1	10.966	10.382	9.457 0.023	9.247 0.023	9.133 0.021
...	4766 790 1	13.208	12.133	10.803 0.030	10.403 0.026	10.284 0.026
HD 290513	4766 828 1	10.852	10.245	8.569 0.019	8.270 0.044	8.239 0.023
HD 290582	4766 811 1	10.864	10.533	10.149 0.026	10.068 0.024	10.026 0.023
HD 290508	4766 767 1	11.220	10.753	9.848 0.023	9.614 0.022	9.581 0.023
HD 290516	4766 756 1	9.447	9.445	9.463 0.022	9.517 0.025	9.500 0.021
...	4766 757 1	12.588	12.425	11.903 0.023	11.792 0.022	11.739 0.021
...	4766 716 1	12.109	11.800	10.653 0.022	10.362 0.023	10.291 0.023
...	4766 698 1	11.433	10.785	9.714 0.022	9.467 0.022	9.389 0.019
HD 290583 A	4766 682 1	11.204	10.792	9.885 0.026	9.739 0.023	9.673 0.026
HD 290515	4766 662 1	9.316	9.246	9.258 0.024	9.257 0.024	9.254 0.023
HD 290507	4766 647 1	11.964	11.337	10.022 0.024	9.752 0.022	9.645 0.023
...	4766 631 1	11.253	10.323	9.072 0.029	8.702 0.038	8.585 0.019
HD 36669	4766 548 1	8.888	8.920	9.004 0.024	9.060 0.026	9.040 0.023
...	4766 528 1	12.330	12.224	11.474 0.050	11.328 0.056	11.256 0.058
...	4766 516 1	14.859	11.151	7.064 0.019	6.090 0.042	5.810 0.034
...	4766 454 1	12.083	11.463	8.997 0.027	8.301 0.067	8.126 0.021
HD 290580	4766 440 1	11.024	10.618	9.811 0.022	9.633 0.024	9.594 0.026
...	4766 877 1	12.312	11.785	10.318 0.023	9.987 0.022	9.965 0.023
HD 36605	4766 881 1	8.050	7.961	7.645 0.020	7.650 0.029	7.636 0.020
HD 290579	4766 889 1	11.425	10.480	9.219 0.023	8.877 0.040	8.814 0.019
...	4766 1062 1	10.820	9.675	7.842 0.030	7.342 0.046	7.205 0.021
HD 290504	4753 653 1	11.210	10.821	9.333 0.026	9.118 0.025	9.031 0.025
...	4766 1071 1	11.707	11.138	10.159 0.022	9.873 0.035	9.817 0.019
...	4766 1087 1	11.386	10.227	8.104 0.027	7.565 0.042	7.466 0.024
...	4766 1103 1	12.627	11.960	9.773 0.028	9.250 0.027	9.096 0.026
HD 290578	4766 1109 1	10.961	10.671	9.994 0.026	9.918 0.022	9.822 0.023
...	4766 1263 1	12.274	11.847	10.926 0.023	10.628 0.026	10.547 0.019
...	4766 1168 1	13.096	12.202	11.294 0.023	10.986 0.024	10.924 0.023
...	4766 1153 1	12.494	11.422	8.974 0.027	8.366 0.046	8.205 0.025
...	4766 1160 1	11.897	11.071	10.174 0.023	9.986 0.024	9.902 0.021
...	4766 994 1	12.405	11.360	9.308 0.024	8.736 0.046	8.595 0.021
HD 290503	4753 832 1	11.662	11.246	10.145 0.024	9.882 0.026	9.789 0.022
...	4766 2124 1	11.996	12.589	9.111 0.043	8.462 0.020	8.351 0.031
HD 290492 AB	4766 915 1	9.447	9.333	8.990 0.030	8.931 0.026	8.919 0.023
...	4766 864 1	12.536	11.511	9.400 0.022	8.844 0.024	8.703 0.023
HD 36760	4766 870 1	7.494	7.569	7.801 0.027	7.861 0.040	7.881 0.026
...	4753 919 1	12.835	11.705	10.953 0.024	10.674 0.024	10.592 0.022
HD 290502	4753 876 1	11.626	11.671	10.584 0.022	10.499 0.022	10.435 0.023
SS 28	4766 937 1	12.038	11.564	9.976 0.026	9.429 0.022	9.244 0.021
HD 36683	4766 956 1	9.825	8.413	6.345 0.026	5.790 0.046	5.634 0.023
HD 290501	4753 808 1	11.275	10.279	8.685 0.018	8.295 0.047	8.202 0.023
BD–00 984	4766 2446 1	8.296	8.367	8.581 0.030	8.673 0.063	8.640 0.021
...	4766 2424 1	11.640	11.352	11.208 0.023	11.051 0.021	11.012 0.023
HD 290493	4766 1041 1	10.129	10.024	9.759 0.024	9.733 0.024	9.732 0.023
HD 290500	4753 595 1	11.468	11.472	10.266 0.024	9.951 0.026	9.587 0.025
HD 36841	4766 1122 1	8.583	8.549	8.432 0.023	8.454 0.031	8.403 0.036
HD 290644	4766 1140 1	9.915	9.347	8.319 0.026	8.051 0.046	8.011 0.017

Table A.2. Photometry of correlated Tycho-2/2MASS sources in the Mintaka field (cont.).

Name	TYC	B_T (mag)	V_T (mag)	$J \pm \delta J$ (mag)	$H \pm \delta H$ (mag)	$K_s \pm \delta K_s$ (mag)
V1093 Ori	4766 1181 1	8.134	8.174	8.279 0.029	8.285 0.042	8.301 0.021
HD 290499	4753 431 1	11.816	11.134	9.889 0.024	9.636 0.026	9.509 0.020
...	4753 415 1	12.200	10.901	8.592 0.026	8.027 0.051	7.800 0.020
HD 290574	4766 1251 1	11.318	10.984	10.179 0.027	10.068 0.026	9.985 0.021
HD 290494	4766 1269 1	10.532	9.822	8.706 0.024	8.417 0.042	8.378 0.023
Mintaka AE–D	4766 2445 1	2.006	2.195	2.744 0.268	2.981 0.226	3.023 0.260
...	4753 49 1	12.106	12.310	11.174 0.024	11.038 0.026	10.994 0.023
HD 36485	4766 2444 1	6.673	6.828	7.165 0.026	7.234 0.027	7.282 0.023
HD 36778	4766 1174 1	9.266	9.261	9.279 0.024	9.300 0.030	9.284 0.021
HD 290495	4753 94 1	10.821	9.224	6.719 0.020	6.072 0.018	5.860 0.023
HD 36694	4766 1146 1	9.427	9.149	8.667 0.030	8.620 0.071	8.577 0.021
HD 290573	4766 1108 1	10.663	10.243	9.408 0.022	9.198 0.030	9.138 0.021
...	4766 2264 1	10.706	10.306	9.580 0.026	9.454 0.022	9.378 0.021
HD 290497	4753 289 1	9.688	9.491	9.009 0.024	8.918 0.025	8.866 0.023
HD 290496	4753 304 1	11.458	10.958	10.059 0.026	9.821 0.024	9.742 0.021
HD 290489	4766 1286 1	10.584	10.172	9.271 0.022	9.062 0.022	8.991 0.019
HD 290568	4766 861 1	11.838	9.771	6.261 0.018	5.453 0.044	5.168 0.018
HD 36726	4766 2300 1	8.937	8.828	8.654 0.024	8.732 0.031	8.674 0.019
...	4766 1320 1	12.199	11.604	10.566 0.026	10.366 0.022	10.260 0.023
...	4766 1335 1	12.564	11.481	10.885 0.024	10.715 0.023	10.639 0.019
HD 290566	4766 1347 1	11.752	11.674	11.220 0.027	11.153 0.022	11.105 0.023
HD 36117	4753 115 1	8.118	7.982	7.717 0.021	7.677 0.047	7.652 0.015
HD 36709	4766 1354 1	8.269	8.295	8.360 0.023	8.387 0.040	8.400 0.023
HD 36139	4753 71 1	6.952	6.879	6.700 0.021	6.707 0.051	6.666 0.018
...	4766 1361 1	12.769	11.612	10.676 0.036	10.278 0.035	10.172 0.027
HD 36840	4766 1369 1	7.518	6.355	5.010 0.240	4.297 0.206	4.263 0.354
...	4766 1380 1	12.364	12.339	10.916 0.024	10.756 0.029	10.700 0.024
...	4766 1382 1	11.791	11.434	10.235 0.026	9.979 0.025	9.900 0.021
HD 36558	114 2260 1	8.237	6.352	3.356 0.222	2.434 0.206	2.241 0.270
...	114 519 1	11.607	11.163	10.211 0.026	9.983 0.022	9.937 0.021
HD 36312	114 471 1	8.058	8.091	8.220 0.019	8.265 0.019	8.277 0.023
HD 290567	114 369 1	11.060	10.529	9.480 0.022	9.196 0.022	9.175 0.021
...	114 216 1	12.324	11.321	9.452 0.026	8.804 0.038	8.762 0.023
...	114 167 1	13.173	11.301	7.723 0.024	6.946 0.047	6.649 0.026
HD 36443	114 64 1	9.238	8.432	7.063 0.021	6.794 0.026	6.656 0.021
...	114 77 1	11.986	11.531	10.077 0.026	9.827 0.023	9.713 0.021
HD 290564	114 171 1	11.518	10.879	10.083 0.021	9.901 0.023	9.872 0.017
...	114 166 1	12.422	12.229	11.277 0.027	11.023 0.024	10.934 0.021
...	114 201 1	13.307	11.816	9.829 0.024	9.349 0.020	9.205 0.020
...	114 203 1	12.499	11.797	10.966 0.024	10.712 0.022	10.672 0.022
...	101 2094 1	12.081	11.525	10.930 0.031	10.756 0.024	10.655 0.024
...	101 2006 1	13.408	11.549	9.826 0.027	9.268 0.023	9.088 0.021
HD 290486 AB	101 1971 1	9.783	8.577	6.586 0.026	6.053 0.044	5.945 0.020
HD 290487	114 359 1	10.316	9.815	8.887 0.021	8.667 0.021	8.619 0.020
...	114 431 1	13.876	11.499	8.550 0.029	7.857 0.044	7.611 0.033
...	114 419 1	11.814	11.149	10.187 0.026	9.958 0.025	9.894 0.023
...	114 479 1	12.640	12.309	10.576 0.018	10.280 0.021	10.190 0.025
...	114 480 1	12.457	11.779	10.672 0.025	10.469 0.023	10.427 0.025
...	114 551 1	12.279	11.807	10.661 0.024	10.442 0.019	10.395 0.023
...	114 535 1	10.716	10.049	8.725 0.023	8.444 0.021	8.383 0.022
HD 290559	114 484 1	11.750	10.724	8.844 0.025	8.340 0.026	8.246 0.023
HD 290484	114 197 1	11.590	11.213	10.660 0.023	10.572 0.024	10.544 0.024
HD 290483	114 439 1	10.395	9.901	8.819 0.029	8.613 0.036	8.572 0.020

Table A.2. Photometry of correlated Tycho-2/2MASS sources in the Mintaka field (cont.).

Name	TYC	B_T (mag)	V_T (mag)	$J \pm \delta J$ (mag)	$H \pm \delta H$ (mag)	$K_s \pm \delta K_s$ (mag)
...	114 263 1	12.232	11.069	8.819 0.029	8.613 0.036	8.572 0.020
...	114 131 1	13.353	11.392	9.271 0.020	8.739 0.018	8.634 0.015
...	114 62 1	12.762	12.236	10.668 0.020	10.413 0.018	10.348 0.015
...	114 288 1	12.616	12.432	11.501 0.027	11.413 0.025	11.360 0.025
...	114 344 1	13.086	12.008	9.344 0.024	8.735 0.023	8.586 0.021
...	114 486 1	12.300	11.534	10.632 0.025	10.381 0.020	10.309 0.022
HD 290560	114 434 1	11.535	11.352	10.257 0.024	10.074 0.021	10.031 0.021

Table A.3. Known young stars in the Alnilam field and in Tycho-2/2MASS.

Name		Sp. type	α (J2000)	δ (J2000)	Features of youth	Reference(s)
HD 36980 AB	*	B9.5V	05 35 15.17	–01 54 07.6	OBA	SCo71, Gu76, Gi83
RX J0535.6–0152 AB	*	G6V	05 35 33.39	–01 52 35.7	T Tau, Li I, X-rays	Al96, Al00, Ma07
HD 290691		B9V	05 35 58.54	–01 51 46.6	OBA	Ne95
HD 290686		A5V	05 36 18.37	–01 50 49.7	OBA	Ne95
HD 290684		A2V	05 35 35.83	–01 46 51.7	OBA	AK59, Gu76, Gi83
HD 37285 AB	*	A0V	05 37 20.41	–01 46 16.3	OBA	Gu76, Gi83
HD 37113		B9.5V	05 35 58.98	–01 45 46.6	OBA	CB66, Gu76, WH78
HD 37389	*	A0V	05 38 08.01	–01 45 07.8	OBA, Vega-like	Gu76, Ou92, BM00
HD 37272		B5V	05 37 14.52	–01 40 03.8	OBA, X-rays	BC72, Gu76, Gr92, *cs08*
HD 37149	*	B6Vnm	05 36 17.83	–01 38 07.2	OBA, Hα	Bi65, Gu76, CBD'I78
HD 290770	*	B8Vne	05 37 02.45	–01 37 21.4	HAeBe, OBA, mIR	Bi65, DHu91, *cs08*
HD 37344	*	B9V	05 37 49.15	–01 35 09.3	OBA	SCh71, Gu76, WH78
HD 290602	*	A3V	05 33 50.05	–01 33 07.9	OBA	Ne95
HD 290678		A0V	05 36 03.07	–01 32 08.9	OBA	Ne95
HD 290677		A5V	05 36 16.38	–01 31 11.8	OBA	Gu76, WH78, GC93
HD 290674	*	A0V	05 36 57.32	–01 26 07.4	OBA	Gu76, Gi83, Ne95
HD 37321 AB	*	B5Vsn	05 37 34.81	–01 25 19.6	OBA	Sh52, AL77, AJ98
HD 36955	*	A0Vp	05 35 04.54	–01 24 06.6	OBA	Gi83, GC93, Ku06
HD 290766		A0V	05 38 21.62	–01 21 51.3	OBA	Ne95
HD 290672		A2Van	05 36 56.70	–01 18 26.6	OBA	WH77, GC93
HD 290765		A0V	05 37 43.90	–01 17 21.4	OBA	Gu76, Gi83
V1247 Ori	*	A5III	05 38 05.25	–01 15 21.7	HAeBe, OBA, Hα, mIR	SCo71, MacC82, *cs08*
Alnilam A	*	B0Iab:	05 36 12.81	–01 12 06.9	OBA, Hα, X-rays	Fr09, MKK43
HD 37397	*	B4Vn	05 38 13.74	–01 10 09.0	OBA	Sh52, AL77
VV Ori AB	*	B1.0V+B4.5V	05 33 31.45	–01 09 21.9	OBA, mIR	Da16, TMS07, *cs08*
HD 36684 AB	*	B9V	05 33 25.20	–01 03 31.4	OBA	CB66, Gu76
HD 290750	*	A5III	05 38 10.87	–01 03 00.9	OBA	SCo71, Gu76
HD 36779 AC	*	B2.5V	05 34 03.89	–01 02 08.6	OBA	Sh52, SCh71
HD 36996		A0V	05 35 21.29	–01 02 00.9	OBA	Gu76, Gi83
HD 37187	*	B9V	05 36 37.07	–01 01 40.8	OBA	Sh52, Gu76
HD 290671	*	B9.5V	05 35 48.90	–00 59 21.2	OBA, X-rays	Gu76, WH78, *cs08*
HD 37076	*	B9V	05 35 48.03	–00 59 12.8	OBA	CB66, Gu76, Gr92
HD 290748		A4V	05 37 28.91	–00 55 47.3	OBA	Gu76, Ne95
HD 37075		A0V	05 35 45.02	–00 54 05.2	OBA	Gu76, Gi83
HD 37302		B9.5V	05 37 36.70	–00 53 13.7	OBA	Gu76, Gi83
HD 290666		A1V	05 34 36.93	–00 53 00.3	OBA	SCo71, Gu76
HD 290665	*	A0Vp	05 35 10.51	–00 50 13.0	OBA	Gu76, Gi83
HD 36915		B8V	05 35 00.30	–00 48 56.0	OBA	AK59, Sh74, Gr92
HD 37112		B6V	05 36 03.58	–00 46 47.8	OBA, X-rays	AK59, CB66, Gu76, *cs08*
V1379 Ori	*	B5V	05 37 45.89	–00 46 41.7	OBA	SCh71, WH78
HD 290662	*	A0Vp(Fe II)	05 35 55.51	–00 46 39.2	OBA, X-rays	SCo71, Gi83, *cs08*
HD 36825		A0V	05 34 24.88	–00 45 47.8	OBA	AK59, Sh74
HD 36954 AB	*	B4Vsn	05 35 12.79	–00 44 07.3	OBA, X-rays	Ne43, Sh52, Me83, *cs08*
HD 37235	*	B8Vp	05 36 57.37	–00 42 06.9	OBA, mIR?	SCh71, BC72, *cs08*
HD 37284		A1III–IV	05 37 23.23	–00 41 50.9	OBA	SCh71, Gi83
HD 290648	*	A2V	05 34 55.28	–00 36 59.1	OBA	Gu76, Gi83, Ne95
HD 290660	*	A3V	05 36 29.75	–00 36 25.0	OBA	Ne95
HD 290745		A2V	05 37 46.94	–00 34 17.1	OBA	Ne95
HD 290650	*	A4Vn	05 35 36.94	–00 30 21.2	OBA	Gu76, Ne95

Table A.4. Known young stars in the Mintaka field and in Tycho-2/2MASS.

Name		Sp. type	α (J2000)	δ (J2000)	Features of youth	Reference(s)
HD 290516		B9.5V	05 30 16.50	–00 53 29.4	OBA	WH78, Gu81
HD 290515	*	A0V	05 30 29.15	–00 50 50.7	OBA	AK59, Gu81
HD 36669		B9V	05 33 25.00	–00 47 01.5	OBA	Gu76, AL77
HD 36605		B8V	05 32 49.65	–00 42 47.2	OBA	CB66, Gu76
HD 290578		B9V	05 32 41.24	–00 38 05.6	OBA	Gu79, Ne95
HD 290492 AB	*	A0.5Vb+	05 31 18.75	–00 29 24.2	OBA	WH78, PG97
HD 36760		B7V	05 34 02.26	–00 28 34.8	OBA	CB66, Gu76
HD 290502		A5V	05 29 31.49	–00 27 28.1	OBA	Gu79, Ne95
SS 28	*	G5:	05 32 43.80	–00 27 15.2	T Tau, Hα, X-rays	SS77, Bo88, Wi89
BD–00 984	*	B8III	05 32 04.00	–00 25 44.8	OBA	WH78, Gu81, WL99
HD 290493		A0V	05 30 21.30	–00 25 13.4	OBA	Gu79, Gu81, Ne95
HD 290500	*	A2V	05 29 48.05	–00 23 43.5	HAeBe, OBA, mIR	Co95, Ne95, Vi03
HD 36841	*	B5Vn	05 34 33.72	–00 23 11.5	OBA	MH55, Gu76
V1093 Ori AB	*	B9Vp(Si)	05 30 45.23	–00 22 24.2	OBA	Gu76, Bo81, No84
HD 290574		B8V	05 33 22.45	–00 20 03.6	OBA	Gu79, Ne95
Mintaka AE–D	*	O9.5II+B0.5II+B:	05 32 00.40	–00 17 56.7	OBA, X-rays	Ha1904, Ha02
Mintaka C	*	B2Vp	05 32 00.41	–00 17 04.4	OBA	KH50, Bo87
HD 36778		B9.5V	05 34 05.86	–00 16 14.6	OBA	AK59, Gu81
HD 290491	*	A0V	05 31 19.00	–00 11 13.3	OBA	WH78, Ne95
HD 290497		A3Va(n)	05 29 14.94	–00 10 11.1	OBA	Gu81, GC93
HD 36726 A	*	A0Vm	05 33 51.73	–00 04 36.3	OBA	AL77, Pa01
HD 36709		B9.5V	05 33 45.45	–00 01 44.1	OBA	AK59, Gu76
HD 290490		A2IV	05 30 51.52	–00 08 09.3	OBA	WH78, Gu81, Høg98
HD 290572	*	B8V?	05 33 54.54	–00 06 21.2	OBA	CP18, Ne95, Høg98
HD 290488		A2V	05 30 16.46	–00 03 03.7	OBA	Gu79, Ne95, Høg98
HD 290570		A2V	05 34 02.33	–00 00 52.8	OBA	Ne95, Høg98
HD 290569	*	A0V?	05 33 06.01	–00 00 02.6	OBA	Gu79, Ne95, Høg98
HD 36312		B8.5V	05 30 48.68	+00 01 42.7	OBA	CB66, Gu81
HD 290564		B5V	05 32 08.74	+00 07 36.9	OBA	Gu79, Ne95

Table A.5. Stars of unknown association membership status in the Alnilam field and in Tycho-2/2MASS.

Name		Sp. type	α (J2000)	δ (J2000)	$\mu_\alpha \cos\delta$ (mas a^{-1})	μ_δ (mas a^{-1})	Remarks	Reference(s)
HD 37038 AB	*	F0V	05 35 32.87	–01 50 46.1	+4.3±1.0	+5.5±1.0		SCo71, No97, No04
TYC 4766 1605 1		–	05 35 08.00	–01 50 19.2	–1.6±1.5	+3.4±1.5		*cs08*
TYC 4767 935 1		–	05 37 32.72	–01 48 23.9	–6.1±2.0	+9.1±2.1		*cs08*
HD 36863	*	A3V	05 34 34.34	–01 44 37.4	–7.5±1.2	–6.8±1.1		Lee68, WH78, Gi83
TYC 4766 1699 1		–	05 36 02.32	–01 44 05.2	+1.9±1.4	–3.9±1.4		*cs08*
TYC 4766 121 1		–	05 34 42.31	–01 40 48.8	+1.6±2.3	–0.6±2.2		*cs08*
HD 290682		F5V	05 35 13.17	–01 40 45.9	+2.7±1.2	+1.7±1.2		Ne95
TYC 4766 1823 1		–	05 36 09.87	–01 40 22.5	+3.0±1.5	+4.0±1.5		*cs08*
TYC 4766 400 1		–	05 34 54.92	–01 38 40.0	–2.4±2.1	–9.5±2.1		*cs08*
TYC 4766 260 1		–	05 34 35.90	–01 33 56.9	–6.3±1.8	–5.8±2.1		*cs08*
TYC 4766 1599 1		–	05 36 59.08	–01 32 31.8	–6.9±1.1	+5.9±1.1		*cs08*
[NYS99] A–06	*	–	05 34 50.54	–01 31 29.8	+2.4±2.1	+0.1±2.1	X-rays	Na99, Ue01, *cs08*
TYC 4766 1833 1		–	05 35 45.00	–01 26 35.5	–5.7±1.5	–3.6±1.7		*cs08*
BD–01 945		A8V	05 33 41.69	–01 25 28.7	–3.0±3.0	–2.6±1.9		Ne95, Høg98
HD 290673	*	A2V	05 36 50.74	–01 25 07.9	+1.8±1.0	–1.0±1.0		Ne95
TYC 4767 1130 1	*	–	05 38 58.63	–01 19 56.7	0	0		*cs08*
HD 290676		F5V	05 35 26.10	–01 19 38.5	–4.6±1.2	+0.0±1.2		Ne95
HD 37172	*	A2Va(n)	05 36 31.60	–01 14 12.2	–5.7±1.1	+0.2±1.1		GC93
TYC 4766 2371 1	*	–	05 35 49.68	–01 14 09.5	–0.7±1.5	+1.9±1.6		*cs08*
TYC 4766 2344 1		–	05 37 05.82	–01 12 54.9	+4.2±1.5	+1.6±1.5		*cs08*
TYC 4766 2221 1		–	05 37 17.40	–01 12 52.2	–0.3±2.0	–2.7±2.1		*cs08*
HD 290670		F8V	05 35 27.54	–01 03 05.1	+1.3±1.2	+0.5±1.2	X-rays	Ne95, *cs08*
TYC 4766 2150 1	*	–	05 37 29.58	–01 02 58.9	+4.0±1.5	–8.0±1.6		*cs08*
TYC 4766 542 1	*	–	05 34 48.70	–01 02 27.6	0	0		*cs08*
HD 290669		F0V	05 34 43.61	–01 00 12.5	–6	+1		Ne95
HD 290753		F8V	05 38 34.62	–00 59 05.3	+1.5±1.1	–1.9±1.1		Ne95
TYC 4766 1756 1		–	05 37 26.49	–00 57 47.4	0	0		*cs08*
HD 290751		F5V	05 38 10.06	–00 55 58.7	+1.9±1.9	+0.3±2.0		Ne95
HD 290664		F5V	05 36 38.14	–00 50 37.0	+3.1±1.2	+1.4±1.2	X-rays	Ne95, *cs08*
HD 290747		F0V	05 37 36.14	–00 49 33.7	–1.9±2.3	–0.2±1.7		Ne95, Høg98
HD 290663		F0V	05 36 27.36	–00 48 25.7	+2.4±1.2	–7.8±1.2	X-rays	Ne95, *cs08*
HD 290661		A9IV	05 35 55.11	–00 42 51.7	0	0		SCo71, Høg98
HD 290746	*	G0V?	05 37 39.75	–00 38 25.9	+0.3±1.9	–2.2±1.9		Ne95
HD 290649		F0V	05 35 16.82	–00 35 22.1	–0.2±1.6	–2.4±1.6		Ne95, Høg98

Table A.6. Stars of unknown association membership status in the Mintaka field and in Tycho-2/2MASS.

Name		Sp. type	α (J2000)	δ (J2000)	$\mu_\alpha \cos\delta$ (mas a^{-1})	μ_δ (mas a^{-1})	Remarks	Reference(s)
HD 290585	*	F1V	05 32 21.79	–01 01 19.2	+0.6±1.4	+1.7±1.3		Ne95, He06
TYC 4766 733 1		–	05 31 52.22	–00 58 37.2	+2.2±2.3	–0.9±2.5		*cs08*
HD 290513	*	F0V?	05 30 43.18	–00 56 58.9	–1.8±1.3	–10.7±1.3		Ne95
TYC 4766 790 1	*	–	05 33 10.86	–00 57 30.7	0	0		*cs08*
HD 290582		F0V	05 33 15.51	–00 55 15.7	–1.0±0.9	+2.2±0.9		Ne95
HD 290508		F5V	05 31 03.79	–00 53 55.1	–1.5±1.1	+2.0±1.1		Ne95
TYC 4766 757 1		–	05 32 24.41	–00 53 21.5	+2.8±1.4	–1.6±1.5		*cs08*
HD 290583 A	*	A7V	05 32 14.80	–00 51 22.7	–1.9±1.1	+2.4±1.1		Gu79
HD 290507	*	A5V?	05 30 47.18	–00 50 21.7	+6.0±1.3	+9.6±1.2		Gu79, Ne95
TYC 4766 528 1	*	–	05 30 04.85	–00 46 17.4	+0.0±1.6	+2.4±1.7		*cs08*
HD 290580		A7V	05 33 01.95	–00 43 55.5	+0.4±1.1	–4.2±1.1		Gu79, Ne95
HD 290504	*	A5V?	05 29 33.35	–00 38 57.9	–0.1±1.3	+3.6±1.3		Gu79, Ne95
TYC 4766 1263 1		–	05 32 03.02	–00 35 17.2	+1.9±1.7	+3.3±1.6	X-rays	*cs08*
TYC 4766 1168 1	*	–	05 31 31.44	–00 33 14.4	0	0		*cs08*
HD 290575		F5V	05 31 54.95	–00 33 09.6	+7.5±1.5	–5.7±1.5		Ne95
TYC 4766 2424 1	*	–	05 31 34.16	–00 25 29.3	–0.3±1.4	+4.6±1.4		*cs08*
TYC 4753 49 1	*	–	05 29 37.04	–00 17 19.8	+1.0±1.8	+3.1±2.0		*cs08*
HD 36694		AVm	05 33 37.97	–00 15 43.2	–5.1±1.2	+10.6±1.2		WH78, Gu81
TYC 4766 1320 1		–	05 31 46.42	–00 04 23.6	+4.9±1.6	–0.9±1.8		*cs08*
TYC 4766 1335 1		–	05 31 03.46	–00 03 32.5	+6.4±1.5	–7.4±1.6		*cs08*
HD 290566		AV	05 31 49.30	–00 02 34.4	–3.1±1.4	–4.0±1.4		Gu79, Ne95
TYC 4766 1380 1		–	05 31 42.73	–00 00 10.2	+5.1±1.6	–1.3±1.8		*cs08*
HD 290565		F2V	05 31 52.77	+00 00 46.1	–3.0±1.3	+1.2±1.3		Ne95
TYC 114 77 1		–	05 33 48.84	+00 06 38.0	–1.3±1.6	–7.0±1.6		*cs08*
TYC 114 203 1		–	05 32 41.40	+00 08 25.2	+1.8±2.5	–10.0±2.5		*cs08*
TYC 101 2094 1		–	05 29 49.98	+00 08 29.6	–0.9±1.3	+0.0±1.3		*cs08*
HD 290487		F2V	05 30 33.56	+00 10 39.7	+3.6±1.1	–6.2±1.2		Ne95
TYC 114 551 1		–	05 32 09.20	+00 14 05.6	–0.1±1.6	–1.5±1.6		*cs08*
HD 290484		F0V	05 30 16.92	+00 18 37.4	+5.5±1.5	+0.4±1.5		Ne95
TYC 114 288 1		–	05 32 21.67	+00 23 34.0	–1.2±1.7	–2.8±1.9		*cs08*
HD 290560		F8V	05 32 57.47	+00 15 37.2	+5.5±1.3	–4.9±1.3		Ne95

Table A.7. Fore- and background stars in the Alnilam field and in Tycho-2/2MASS.

Name		Sp. type	α (J2000)	δ (J2000)	$\mu_\alpha \cos\delta$ (mas a^{-1})	μ_δ (mas a^{-1})	Reference(s)
HD 290690		G0V	05 35 44.98	–01 56 03.8	–4.9±1.3	+3.9±1.3	Ne95
TYC 4770 1072 1		K–M:	05 36 26.83	–01 56 03.1	–1.5±1.3	+9.6±1.3	*cs08*
TYC 4766 2029 1		F–G:	05 34 48.41	–01 51 46.8	–11.1±1.8	+8.3±1.8	*cs08*
TYC 4767 3 1		K–M:	05 37 38.35	–01 51 22.9	+0.9±1.3	–9.4±1.3	Kr03, *cs08*
TYC 4766 1891 1		F–G:	05 36 41.21	–01 46 34.2	–6.0±2.0	–33.5±2.1	*cs08*
HD 290685		K0V	05 36 28.07	–01 45 15.5	–9.9±1.0	+3.3±1.1	Ne95
HD 290683		F8V	05 35 51.32	–01 42 20.7	–10.6±1.5	+26.7±1.5	Ne95
IRAS 05354–0142	*	C/S?	05 37 56.58	–01 40 50.2	+1.2±2.0	+5.8±2.0	Kr03, *cs08*
TYC 4766 1643 1		K–M:	05 35 50.47	–01 38 12.0	–3.9±1.4	–19.8±1.4	*cs08*
HD 290681		F2V	05 35 30.45	–01 35 52.8	–14.2±1.0	+2.8±1.1	Ne95
TYC 4766 2063 1		K–M:	05 36 56.83	–01 35 21.0	0	0	*cs08*
HD 290679	*	K2V	05 35 36.39	–01 34 32.7	+2.2±1.2	+3.3±1.3	Ne95
HD 290680	*	G0V?	05 35 26.04	–01 34 29.9	–9.6±1.3	–1.0±1.3	Ne95
TYC 4766 265 1		K–M:	05 35 15.13	–01 33 56.3	+1.3±2.1	–11.4±2.1	*cs08*
TYC 4767 498 1		K–M:	05 38 28.32	–01 32 08.6	–0.7±2.0	–4.4±2.0	*cs08*
HD 36780	*	K0V	05 34 04.05	–01 28 12.9	–11.7±0.9	–23.1±1.0	Gr72, BLM88, Kr03
TYC 4766 144 1		F–G:	05 34 57.10	–01 28 11.6	+2.2±1.6	+19.4±1.6	*cs08*
TYC 4766 2326 1		K–M:	05 35 52.85	–01 25 21.1	–4.8±1.3	–2.8±1.3	*cs08*
HD 290675	*	F4V	05 36 00.02	–01 23 52.4	+8.7±1.1	+20.4±1.1	Gi83, Ne95
HD 290767		K0V	05 38 41.39	–01 23 50.5	+1.3±1.3	–2.6±1.3	Ne95
HD 37506		A0–F1Vm	05 39 04.69	–01 13 50.9	–28.9±1.0	+5.5±1.0	Ba74, Gi83, Re88
TYC 4766 724 1		F–G:	05 33 34.41	–01 12 47.4	+13.5±1.3	–26.0±1.3	*cs08*
HD 290588		K0V	05 33 41.69	–01 11 36.5	+3	+1	Ne95, Kr03
TYC 4767 487 1		K–M:	05 37 42.96	–01 11 32.7	0	0	*cs08*
HD 290762		F8V	05 38 50.25	–01 10 22.0	–19.3±2.3	–1.7±1.7	Ne95, Høg98
HD 37491	*	K0V	05 38 55.29	–01 10 12.7	+33.0±1.0	–14.6±1.0	CP18, Kr03, Ál04
HD 290763		G0V	05 38 23.40	–01 08 20.0	+9.3±1.3	+12.1±1.3	Ne95
TYC 4766 477 1		F–G:	05 33 27.85	–01 03 48.0	–23.2±1.3	+2.3±1.3	*cs08*
HD 37443		F5V	05 38 32.64	–01 03 30.6	–30.3±1.1	+5.8±1.1	Gi83, No04, Kr03
HD 290749	*	B8V?	05 37 30.09	–01 03 25.4	–5.4±1.3	+12.7±1.3	Ne95
HD 36882	*	G5V	05 34 48.19	–00 57 13.1	–12.3±0.9	+37.0±0.9	Eg85, Kr03
TYC 4767 760 1		K–M:	05 38 11.01	–00 56 11.9	+3.9±1.9	–68.5±2.0	*cs08*
HD 290667	*	F7V	05 34 24.23	–00 55 19.1	–1.3±1.9	–27.5±1.7	St86, DK88, MMB03
TYC 4767 1008 1		K–M:	05 38 36.72	–00 50 48.4	0	0	*cs08*
HD 290752		G0V	05 38 02.40	–00 48 42.8	+6.6±1.8	–11.4±1.8	Ne95
StHA 46	*	G6	05 35 55.11	–00 42 51.7	0	0	St86, MMB03
TYC 4767 715 1		K–M:	05 38 15.57	–00 42 25.0	0	0	*cs08*
HD 290647	*	G5V	05 34 28.60	–00 42 20.9	–4	+12	Ne95
TYC 4767 2257 1	*	MIII?	05 36 49.68	–00 37 15.0	0	0	*cs08*

Table A.8. Fore- and background stars in the Mintaka field and in Tycho-2/2MASS.

Name		Sp. type	α (J2000)	δ (J2000)	$\mu_\alpha \cos\delta$ (mas a^{-1})	μ_δ (mas a^{-1})	Reference(s)
HD 290586		G5V	05 32 09.22	–01 01 31.5	+1.9±1.3	–45.9±1.3	Ne95
TYC 4766 644 1		K–M:	05 32 29.78	–01 00 38.8	–2.7±1.5	+1.2±1.5	*cs08*
TYC 4766 687 1		K–M:	05 31 35.28	–00 59 42.9	+2.7±1.5	+1.5±1.5	*cs08*
HD 290509		F0V	05 31 26.41	–00 58 33.1	+29.4±1.1	–9.5±1.1	Ne95
TYC 4766 716 1		F–G:	05 33 09.06	–00 52 30.8	+8.5±1.3	–37.3±1.4	*cs08*
HD 290514		F8V	05 30 38.34	–00 52 09.0	+13.0±1.2	–13.6±1.2	Ne95
HD 290581		G0V	05 33 18.73	–00 50 12.4	+57.0±2.1	–57.8±1.9	Ne95, Høg98
HD 290584		G5V	05 31 54.41	–00 49 56.0	+28.8±1.3	–111.3±1.3	Ne95
TYC 4766 516 1	*	MIII:	05 31 27.11	–00 45 41.3	–0.6±2.1	+5.4±2.1	*cs08*
TYC 4766 454 1		K–M:	05 33 36.55	–00 44 21.9	0	0	*cs08*
TYC 4766 877 1		F–G:	05 32 22.53	–00 42 57.3	+9.1±1.5	–18.4±1.5	*cs08*
HD 290579		F5V	05 32 24.85	–00 42 38.9	+14.2±1.3	–42.9±1.3	Ne95
HD 290647	*	G5V	05 34 28.60	–00 42 20.9	–4	+12	Ne95
HD 290577		G5V	05 32 10.58	–00 39 14.4	+12.9±1.1	–19.5±1.1	Ne95
HD 290646		F5V	05 34 08.88	–00 38 48.7	+12.9±1.4	–35.6±1.4	Ne95
HD 290576	*	K4V	05 32 09.06	–00 38 36.7	–33.7±1.3	–98.5±1.3	St86, Ne95
TYC 4766 1103 1		K:V	05 32 36.20	–00 38 09.1	–62.2±1.6	–119.3±1.7	*cs08*
TYC 4766 1153 1		K–M:	05 30 46.79	–00 33 10.8	+5.4±2.1	–5.3±2.1	*cs08*
TYC 4766 994 1		K–M:	05 32 27.24	–00 30 24.6	–1.8±1.6	+4.0±1.7	*cs08*
HD 290503		G0V	05 29 25.25	–00 30 13.3	+7.3±1.4	+5.0±1.4	Ne95
TYC 4766 2124 1	*	G8III	05 31 17.17	–00 29 30.4	+0.0±1.6	–18.1±1.7	Pa01a
TYC 4766 864 1		K–M:	05 30 12.01	–00 28 35.1	–1.5±1.6	–2.5±1.6	*cs08*
TYC 4753 919 1		F–G:	05 29 33.41	–00 27 57.5	+12.9±2.1	–40.9±2.2	*cs08*
HD 36683		K0V	05 33 27.54	–00 26 28.9	+3.9±1.0	–5.1±1.0	CP18
HD 290501		G5V	05 29 59.96	–00 26 19.1	+25.5±1.3	–6.1±1.3	Ne95
HD 290645		G5V	05 34 08.42	–00 24 03.4	–22.3±2.0	–1.9±1.6	Ne95, Høg98
HD 290644		F8V	05 34 26.36	–00 22 54.7	+37.5±1.0	–61.9±1.0	Ne95
HD 290499		G0V	05 29 30.47	–00 20 44.8	+1.2±1.4	–12.5±1.4	Ne95
HD 290498		K:V	05 29 09.60	–00 20 33.2	+35.8±1.4	–76.4±1.4	Ne95
HD 290494		F8V	05 30 29.55	–00 19 22.0	+19.8±1.1	+15.7±1.1	Ne95
HD 290495		K2V	05 29 58.57	–00 16 01.3	+1.0±1.1	–0.1±1.1	Ne95
HD 290573		F5V	05 33 56.24	–00 14 43.2	–1.8±1.3	–24.5±1.3	Ne95
HD 290643		F5V	05 34 52.98	–00 10 35.9	+11.0±1.8	–18.5±1.7	Ne95, Høg98
HD 290496		G0V	05 29 53.49	–00 09 30.4	–4.0±1.4	–14.6±1.4	Ne95
HD 290489		F5V	05 30 43.83	–00 07 22.8	+19.6±1.2	–0.2±1.2	Ne95
HD 290568	*	M0V	05 32 52.50	–00 07 18.2	–2.5±1.0	–2.4±1.1	Ne95
HD 36117	*	A2Van	05 29 27.38	–00 02 32.2	+12.9±1.1	–15.9±1.2	Gu81, GC93
HD 36139	*	A2V	05 29 37.56	–00 01 16.5	–3.0±1.1	–14.8±1.1	Gu81, ML91
TYC 4766 1361 1		F–G:	05 31 48.79	–00 01 13.7	+3.8±1.6	+27.8±1.8	*cs08*
TYC 4753 70 1		K–M:	05 29 33.32	–00 01 11.5	0	0	*cs08*
HD 36840	*	G5V	05 34 29.28	–00 00 44.4	–6.0±1.1	–4.3±1.1	Eg85
TYC 4766 1382 1		F–G:	05 30 46.38	–00 00 07.1	+3.1±1.6	+31.5±1.6	*cs08*
HD 36558	*	K5V	05 32 37.98	+00 00 43.0	+22.6±1.0	–10.3±1.0	No04
HD 290567		G0V	05 32 30.06	+00 03 00.9	+11.8±1.1	–5.3±1.1	Ne95
TYC 114 216 1		K:	05 32 03.06	+00 04 29.2	+8.2±1.3	–18.2±1.3	*cs08*
TYC 114 167 1		M:	05 31 29.98	+00 04 54.7	0	0	*cs08*
HD 36443	*	G5V	05 31 44.35	+00 05 55.7	+182.1±1.0	–453.6±0.9	Ro55
TYC 114 166 1		F–G:	05 34 10.86	+00 07 46.5	–20.1±1.8	–2.5±1.9	*cs08*
TYC 114 201 1		K–M:	05 32 44.80	+00 08 23.6	0	0	*cs08*
TYC 101 2006 1		K–M:	05 29 59.53	+00 09 56.4	0	0	*cs08*
HD 290486 AB	*	K0V	05 29 45.16	+00 10 34.3	–8.2±1.2	–28.8±1.1	Ne95
TYC 114 431 1		K–M:	05 30 24.87	+00 11 38.8	0	0	*cs08*

Table A.8. Fore- and background stars in the Mintaka field and in Tycho-2/2MASS (cont.).

Name	Sp. type	α (J2000)	δ (J2000)	$\mu_\alpha \cos\delta$ (mas a^{-1})	μ_δ (mas a^{-1})	Reference(s)
HD 290561	G:V	05 32 51.66	+00 11 45.8	–14.8±1.3	–12.5±1.3	Ne95
TYC 114 479 1	F–G:	05 31 04.15	+00 12 22.5	+11.4±2.4	–15.8±2.7	*cs08*
TYC 114 480 1	F–G:	05 31 37.55	+00 12 25.2	+6.9±1.7	–11.1±1.6	*cs08*
HD 290563	G5V	05 31 46.80	+00 14 39.1	–2.2±1.3	–38.0±1.3	Ne95
HD 290559	G5V	05 33 16.98	+00 15 13.4	+10.6±1.3	+12.5±1.3	Ne95
TYC 114 263 1	K–M:	05 30 54.88	+00 17 50.2	+2.7±1.5	–3.5±1.6	*cs08*
TYC 114 131 1	K–M:	05 31 08.49	+00 19 29.4	+2.4±2.1	+1.9±2.1	*cs08*
TYC 114 62 1	G–K:	05 31 19.27	+00 20 19.9	+2.8±1.7	–28.7±1.8	*cs08*
HD 290483	G0V	05 30 45.09	+00 21 26.6	+9.3±1.3	–22.7±1.3	Ne95
HD 290558	F5V	05 33 20.35	+00 21 52.3	–9.6±1.8	–10.5±1.8	Ne95
TYC 114 344 1	K–M:	05 32 45.48	+00 23 21.7	0	0	*cs08*

Table A.9. Known and new young stars in the Alnilam field and in DENIS/2MASS.

Name		α (J2000)	δ (J2000)	$i \pm \delta i$ (mag)	$K_s \pm \delta K_s$ (mag)	Features of youth	Reference(s)
E Ori 2–1328	*	05 36 26.09	–01 39 46.8	16.380 1.00	13.792 0.040	Li I, low *g*	Bé03a
E Ori 2–1868	*	05 36 21.56	–01 37 49.4	18.0 0.3	15.266 0.115	low *g*	Bé03b
E Ori 1–388	*	05 37 54.30	–01 37 20.6	17.397 1.00	13.908 0.049	low *g*	Bé03a
E Ori 1–1644	*	05 36 57.12	–01 36 29.4	16.752 1.00	13.884 0.044	Li I	Bé03a
E Ori 2–878	*	05 36 18.28	–01 27 08.3	17.353 1.00	14.567 0.066	low *g*	Bé03b
E Ori 2–705	*	05 35 59.47	–01 25 38.9	16.8 0.3	14.336 0.059	low *g*	Bé03b
E Ori 2–603	*	05 36 28.57	–01 24 41.0	16.079 1.00	13.234 0.033	low *g*	Bé03b
Kiso A–0904 35		05 35 28.54	–01 25 31.5	14.102 0.03	11.559 0.026	Hα	Wi89
Kiso A–0904 42	*	05 35 54.23	–01 23 05.8	13.1 0.3	11.740 0.018	Hα	Wi89
Kiso A–0904 41	*	05 35 53.49	–01 23 04.4	13.6 0.3	11.431 0.024	Hα	Wi89
2E 1432		05 36 52.31	–01 21 24.7	12.671 1.00	10.686 0.021	X-rays	McD94, *cs08*
V469 Ori	*	05 34 25.82	–01 21 06.5	11.771 0.03	8.759 0.021	Hα, X-rays	HM53, Wi89, Na99
Kiso A–0904 76	*	05 39 02.57	–01 20 32.3	12.277 0.03	9.827 0.023	Li I, Hα	Wi89, Br05
Haro 5–80	*	05 38 04.77	–01 17 18.8	14.624 1.00	11.805 0.025	Hα	HM53, Wi89
2E 1398	*	05 35 55.44	–01 13 31.9	15.3 0.3	10.600 0.021	X-rays	McD94, *cs08*
Kiso A–0904 37	*	05 35 32.15	–01 12 03.6	15.081 0.04	13.899 0.066	Hα	Wi89
StHA 47	*	05 35 22.93	–01 11 24.3	11.415 0.02	9.181 0.023	Hα	St86, DK88
CVSO 162	*	05 38 30.85	–01 10 24.3	13.797 0.03	11.692 0.021	Li I, Hα	Br05, McG06
Kiso A–0904 21		05 34 17.14	–01 09 34.9	13.719 0.03	11.109 0.022	Hα	Wi89
Annizam 363062	*	05 36 34.11	–01 09 16.1	10.202 1.00	9.464 0.021	X-rays	*cs08*
Haro 5–67	*	05 33 43.26	–01 08 34.7	12.000 0.03	9.495 0.018	Hα, Ca II, mIR	HM53, Sa71
Kiso A–0904 50	*	05 36 40.51	–01 07 51.8	14.779 1.00	13.218 0.032	Hα	Wi89
Haro 5–76		05 37 15.45	–01 07 47.7	13.343 1.00	10.992 0.021	Hα	HM53, Wi89
Kiso A–0904 61	*	05 37 31.49	–01 03 33.8	14.727 1.00	11.786 0.027	Hα	Wi89
V583 Ori	*	05 36 35.15	–01 02 16.8	13.907 1.00	10.881 0.019	Hα	HM53, Wi89
HD 36779 B	*	05 34 02.06	–01 02 11.1	9.831 0.03	8.004 0.031	Li I	MMR92, PPR92
Kiso A–0904 20		05 34 15.54	–01 01 03.0	11.957 0.03	10.130 0.019	Hα	Wi89
2E 1357	*	05 35 15.69	–00 59 28.7	12.819 0.02	10.925 0.021	Li I, Hα, X-rays	Al96, Al00
Kiso A–0904 28	*	05 34 54.92	–00 57 30.8	10.794 0.03	8.666 0.024	Hα, X-rays	Wi89, Kr03, *cs08*
Kiso A–0904 30	*	05 35 12.77	–00 54 52.8	13.038 0.02	11.394 0.019	Hα	Wi89
2E 1449	*	05 37 33.78	–00 53 51.8	10.592 1.00	8.923 0.019	Li I, X-rays	Al96, Al00
Kiso A–0904 60	*	05 37 30.04	–00 48 52.1	12.608 1.00	10.486 0.033	Li I, Hα	Wi89, Br05
Kiso A–0904 65	*	05 37 57.96	–00 47 23.6	15.830 1.00	13.070 0.032	Li I, Hα	Wi89, Br05
CVSO 124	*	05 33 49.04	–00 46 26.0	13.479 0.03	11.119 0.021	Li I, Hα	Br05, McG06
Kiso A–0904 34	*	05 35 25.22	–00 43 24.3	14.681 0.03	11.101 0.019	Hα	Wi89, SE05
Kiso A–0904 33	*	05 35 23.63	–00 43 08.8	12.372 0.02	10.111 0.032	Hα	Wi89
V483 Ori		05 35 20.17	–00 42 13.7	12.339 0.02	10.139 0.021	Hα, X-rays	HM53, Wi89, Al96
PU Ori	*	05 36 24.29	–00 42 12.0	12.031 1.00	8.764 0.019	Hα, mIR, [O I]	HM53, HR72
V472 Ori		05 34 48.91	–00 37 16.7	12.188 0.03	9.538 0.019	Hα	HM53, Wi89
StHA 48	*	05 35 43.27	–00 34 36.7	10.983 0.02	9.061 0.019	Li I, Hα	St86, MMB03

Table A.10. Known and new young stars in the Mintaka field and in DENIS/2MASS.

Name		α (J2000)	δ (J2000)	$i \pm \delta i$ (mag)	$K_s \pm \delta K_s$ (mag)	Features of youth	Reference(s)
CVSO 124	*	05 33 49.04	–00 46 26.0	13.479 0.03	11.119 0.021	Li I, Hα	Br05, McG06
Kiso A–0903 231		05 32 29.42	–00 41 39.7	13.209 0.03	11.329 0.021	Hα	Ko89
Kiso A–0903 221	*	05 32 10.16	–00 37 12.4	12.488 0.03	10.662 0.021	Hα	Ko89
Kiso A–0904 4	*	05 32 25.78	–00 36 53.3	12.895 0.03	10.287 0.021	Li I, Hα	Wi89, Ko89, Br05
Kiso A–0904 10		05 33 00.78	–00 36 20.8	11.271 0.03	9.170 0.021	Hα	Ko89
IRAS 05307–0038	*	05 33 18.68	–00 36 14.1	11.859 0.03	8.627 0.021	Li I, Hα, Hβ, mIR	YC94, Yun97
Kiso A–0903 240		05 32 50.45	–00 35 42.3	14.206 0.04	11.753 0.027	Hα	Ko89
Kiso A–0904 22	*	05 34 26.36	–00 33 51.2	14.424 0.04	11.880 0.019	Hα	Ko89
Kiso A–0904 13	*	05 33 06.56	–00 22 54.7	13.409 0.03	10.700 0.021	Hα	Wi89, Ko89
Mantaqah 487126	*	05 32 27.11	–00 22 46.0	13.663 0.03	11.469 0.021	X-rays	*cs08*
1AXG J053127–0021	*	05 31 27.77	–00 20 48.1	10.8 0.3	10.391 0.021	X-rays	Ue01, *cs08*
Kiso A–0903 185		05 30 01.34	–00 20 45.8	13.914 1.00	11.256 0.027	Hα	Ko89
Mantaqah 148186	*	05 31 59.24	–00 20 26.2	15.309 0.04	12.638 0.030	X-rays	*cs08*
Mantaqah 400105	*	05 32 26.37	–00 19 39.2	13.674 0.03	11.402 0.019	X-rays	*cs08*
Kiso A–0904 29		05 34 59.57	–00 18 59.8	13.821 0.03	11.302 0.023	Hα	Wi89
Mantaqah 320042	*	05 32 14.68	–00 13 55.8	13.540 0.03	11.244 0.023	X-rays	*cs08*
Kiso A–0904 18	*	05 33 59.85	–00 05 06.5	13.892 0.04	10.717 0.026	Hα	Wi89
Kiso A–0903 227		05 32 25.13	–00 04 36.3	13.984 0.04	11.783 0.023	Hα	Ko89

Table A.11. Stars of unknown status in the Alnilam and Mintaka fields and in DENIS/2MASS.

Name		α (J2000)	δ (J2000)	$i \pm \delta i$ (mag)	$K_s \pm \delta K_s$ (mag)	Remarks	Reference(s)	Field
E Ori 1–726		05 37 39.00	–01 39 46.8	17.966 1.00	15.035 0.150	M6.0	Bé03a	Alnilam
E Ori 2–1982	*	05 36 05.54	–01 39 00.5	19.0 0.3	15.152 0.110	M8.0:	Bé03b	Alnilam
[NYS99] B–16		05 38 35.98	–01 38 31.6	15.426 0.05	13.532 0.034	X-rays?	Na99	Alnilam
[OSP2002] OriI–2N 4	*	05 38 02.60	–01 34 39.3	16.560 1.00	14.096 0.059	Hα?	OGP02	Alnilam
2E 1343		05 34 58.20	–01 12 37.9	11.635 0.02	10.296 0.021	X-rays?	McD94	Alnilam
BD–01 969B	*	05 36 22.96	–01 10 30.9	9.968 1.00	7.643 0.027	"Alnilam B"	Sc1886	Alnilam
Kiso A–0903 241		05 32 51.09	–01 00 07.1	14.761 0.05	13.597 0.048	mid/faint Hα, blue $I - K_s$	Ko89	Mintaka
AG–00 669	*	05 35 11.50	–00 54 30.5	10.674 0.02	10.257 0.019	early A-type?	Ko69	Alnilam
Kiso A–0903 239		05 32 50.10	–00 49 43.3	13.912 0.04	12.726 0.026	faint Hα, blue $I - K_s$	Ko89	Mintaka
Kiso A–0903 222		05 32 15.69	–00 47 43.4	15.607 0.05	14.260 0.077	mid/faint Hα, blue $I - K_s$	Ko89	Mintaka
Kiso A–0903 237		05 32 45.53	–00 37 48.4	14.075 0.04	12.774 0.034	mid/faint Hα, blue $I - K_s$	Ko89	Mintaka
Kiso A–0903 183		05 29 53.05	–00 20 12.0	13.249 1.00	12.201 0.026	faint Hα, blue $I - K_s$	Ko89	Mintaka
Kiso A–0904 19		05 34 14.87	–00 19 50.6	14.094 0.04	13.133 0.028	faint (?) Hα, blue $I - K_s$	Wi89	Mintaka
BD–00 983B	*	05 31 58.75	–00 18 18.7	12.253 0.03	11.223 0.021	"Mintaka B"	Sc1886	Mintaka
BD–00 993B	*	05 33 50.99	–00 04 53.0	10.867 0.03	9.903 0.023	"HD 36726 BC"	Sc1886	Mintaka
Kiso A–0904 7		05 32 50.51	–00 01 56.7	14.987 0.05	13.614 0.039	very faint Hα, blue $I - K_s$	Wi89	Mintaka
Kiso A–0904 11		05 33 05.71	+00 12 29.1	15.528 0.06	14.488 0.088	faint Hα, blue $I - K_s$	Wi89	Mintaka

Table A.12. Fore- and background objects in the Alnilam and Mintaka fields and in DENIS/2MASS.

Name		α (J2000)	δ (J2000)	$i \pm \delta i$ (mag)	$K_s \pm \delta K_s$ (mag)	Sp. type	References	Field
X Ori	*	05 37 38.65	–01 46 16.7	11.4–15.5 (V_T)	1.55 0.05	M8	Wo1904, TMW05	Alnilam
HD 290769		05 37 52.38	–01 27 37.2	10.25 1.0	9.735 0.023	F8	Ne95	Alnilam
Ruber 1	*	05 31 51.40	–00 45 51.2	15.717 0.05	12.927 0.027	late MV:	*cs08*	Mintaka
G 99–18	*	05 34 00.01	–00 42 03.5	11.941 0.03	10.056 0.019	K–M:	GBT61, SG03	Aln.+Min.
HD 290571		05 34 00.31	–00 03 49.6	10.634 0.03	9.989 0.023	F9	Ne95	Mintaka
HD 290562		05 32 39.45	+00 11 30.3	11.068 0.04	10.433 0.022	F5	Ne95	Mintaka
Ruber 2	*	05 33 11.20	+00 13 58.4	15.346 0.06	12.144 0.028	late MIII:	*cs08*	Mintaka

Table A.13. Probable reddened stars in DENIS/2MASS associated to remnant molecular clouds in the Alnilam and Mintaka fields.

2MASS designation	$i \pm \delta i$ (mag)	$J \pm \delta J$ (mag)	$H \pm \delta H$ (mag)	$K_s \pm \delta K_s$ (mag)	Associated to cloud	Field
2MASS J05380477–0145322	16.743 1.00	13.557 0.042	12.735 0.035	12.164 0.030	ORI I–2	Alnilam
2MASS J05332555–0038348	16.691 0.10	14.120 0.030	12.799 0.032	12.285 0.026	IC 423	Mintaka
2MASS J05332310–0038196	18.251 0.21	15.961 0.076	14.853 0.055	14.291 0.077	IC 423	Mintaka
2MASS J05332631–0037462	16.335 0.08	14.272 0.033	13.057 0.029	12.590 0.027	IC 423	Mintaka
2MASS J05332263–0036565	18.258 0.21	15.425 0.049	14.072 0.032	13.397 0.029	IC 423	Mintaka
2MASS J05332198–0036306	16.912 0.11	14.896 0.033	13.757 0.027	13.174 0.021	IC 423	Mintaka
2MASS J05332038–0036225	16.251 0.08	14.415 0.026	13.489 0.022	13.065 0.036	IC 423	Mintaka

Table A.14. Galaxies in the Alnilam field and in 2MASS.

Name		Name		Name	
2MASX J05355304–0156495		2MASX J05363979–0156145		2MASX J05362575–0154324	
2MASX J05364032–0152595		2MASX J05350166–0152546		2MASX J05360480–0151535	
2MASX J05360091–0151165		2MASX J05365752–0150543		2MASX J05374550–0150164	
2MASX J05375584–0146454		2MASX J05344538–0145356		2MASX J05381044–0145266	
2MASX J05371064–0145203		2MASX J05352182–0144308		2MASX J05371671–0144223	
2MASX J05365111–0143514		2MASS J05345451–0143256	*	2MASX J05372418–0141053	
2MASX J05381458–0139166		2MASX J05365376–0138594		2MASX J05350345–0138506	
2MASS J05352819–0138211		2MASX J05383539–0138065		2MASX J05382719–0137545	
2MASX J05383626–0137415		2MASX J05373249–0137354		2MASX J05361998–0136194	
2MASX J05383159–0136005		2MASX J05375063–0133364		2MASS J05382417–0132226	
2MASS J05340366–0131239		2MASX J05352108–0131218		2MASX J05380604–0131146	
2MASS J05372319–0130518		2MASX J05380691–0130536		2MASX J05383486–0130375	
2MASX J05364774–0129534		2MASX J05340830–0129455		2MASS J05372731–0128272	
2MASX J05350452–0128016		2MASS J05341517–0126479		2MASX J05370910–0123053	
2MASX J05370124–0122223		2MASX J05370444–0119143		2MASX J05370030–0119123	
2MASX J05364552–0118064		2MASX J05370270–0118043		2MASX J05364771–0117164	
2MASX J05334401–0115292		2MASX J05343097–0110405		2MASX J05331784–0107542	
2MASX J05354247–0107405		2MASX J05340262–0107362		2MASX J05333091–0106382	
2MASX J05332498–0106242	*	2MASX J05355727–0106195		2MASX J05374383–0105424	
2MASX J05333921–0105282		2MASX J05360287–0105154		2MASX J05385334–0104469	
2MASX J05353520–0103285		2MASX J05333841–0101242		2MASX J05354333–0059365	
2MASX J05371457–0058193		2MASX J05390342–0057599		2MASX J05363011–0056183	
2MASX J05335414–0052272		2MASX J05361037–0051135		2MASS J05353734–0046354	
2MASX J05372731–0046063		2MASX J05342997–0045145		2MASX J05362823–0044113	
2MASX J05351374–0043198		2MASX J05343902–0042576		2MASX J05365804–0042413	*
2MASX J05343722–0041396		2MASX J05343163–0041185		2MASX J05345216–0041156	
2MASX J05365250–0040504		2MASX J05345349–0040316		2MASX J05360408–0039455	
2MASX J05364723–0039144	*	2MASX J05365543–0039133		2MASS J05364746–0039110	*
2MASS J05365167–0032170					

Table A.15. Galaxies in the Mintaka field and in 2MASS.

Name		Name	Name	
2MASX J05325501–0053516		2MASX J05314416–0053202	2MASX J05305116–0052384	
2MASX J05335414–0052272		2MASX J05314556–0050562	2MASS J05315182–0050178	
2MASS J05314896–0049143		2MASS J05314022–0049037	2MASX J05323724–0048535	
2MASX J05325934–0048476		2MASX J05323377–0048285	2MASX J05325308–0048086	
2MASX J05313329–0047492		2MASX J05320003–0045342	2MASX J05321351–0045065	
2MASX J05335608–0042102		2MASX J05343163–0041185	2MASS J05330941–0040445	
2MASX J05330015–0039056		2MASX J05301419–0037282	2MASX J05314250–0028332	
2MASX J05293010–0026402		2MASX J05340527–0025102	2MASS J05330277–0025041	
2MASX J05341703–0024415		2MASX J05320297–0023572	2MASX J05343129–0022225	
2MASX J05332567–0021282		2MASX J05323238–0021065	2MASX J05321205–0020595	
2MASS J05291137–0020560		2MASX J05291407–0020349	2MASX J05324032–0016435	
2MASX J05301773–0013502		2MASX J05311710–0013482	2MASS J05294580–0012150	
2MASX J05292344–0008232		2MASX J05322692–0007285	2MASX J05322252–0006505	
2MASX J05314258–0006312		2MASS J05322588–0005429	2MASS J05341337–0044087	*
2MASS J05313608–0008425		2MASX J05314865–0003032	2MASX J05341949–0002255	
2MASX J05335580–0001192		2MASX J05324949–0001066	2MASX J05314030–0001022	
2MASX J05322266–0000555	*	2MASX J05320791–0000472	2MASX J05325495+0000443	
2MASX J05341820+0001179		2MASX J05342840+0002389	2MASX J05295174+0004467	
2MASX J05313717+0005107		2MASX J05335592+0005426	2MASX J05312760+0006160	
2MASX J05332801+0006313		2MASX J05335178+0006456	2MASX J05332467+0006563	
2MASX J05333141+0007493		2MASX J05315248+0008553	2MASS J05333785+0009141	
2MASX J05305539+0011132		2MASX J05313940+0014010	2MASX J05304639+0017192	
2MASX J05312213+0018280				

Table A.16. Young star and brown dwarf candidates in the Alnilam field and in DENIS/2MASS.

Identification		α (J2000)	δ (J2000)	$i \pm \delta i$ (mag)	$J \pm \delta J$ (mag)	$H \pm \delta H$ (mag)	$K_s \pm \delta K_s$ (mag)
Annizam 2511174		05 36 30.72	–01 53 43.9	14.669 1.00	13.059 0.026	12.321 0.032	12.103 0.026
Annizam 2445191		05 35 42.05	–01 52 08.2	16.014 0.06	14.264 0.031	13.629 0.040	13.279 0.032
Annizam 2557157		05 37 19.82	–01 51 18.7	14.686 1.00	13.116 0.023	12.483 0.028	12.176 0.021
Annizam 2609154		05 37 28.93	–01 51 13.2	17.777 1.00	15.178 0.039	14.567 0.035	14.163 0.064
Annizam 2511204		05 35 05.80	–01 50 28.3	16.424 0.07	14.597 0.035	13.960 0.043	13.726 0.039
Annizam 2306164		05 36 55.48	–01 49 02.0	16.444 1.00	14.739 0.030	14.133 0.038	13.746 0.035
Annizam 2292164		05 36 54.01	–01 48 54.2	16.538 1.00	14.459 0.028	13.773 0.038	13.505 0.035
Annizam 2684144		05 37 57.03	–01 48 29.8	16.584 1.00	14.603 0.035	13.962 0.032	13.723 0.054
Annizam 2206168		05 36 44.28	–01 48 02.2	14.782 1.00	13.050 0.027	12.429 0.037	12.175 0.023
Annizam 2692142		05 38 02.32	–01 47 40.5	16.488 1.00	14.088 0.030	13.487 0.033	13.103 0.034
Annizam 2382152		05 37 26.75	–01 47 15.2	16.312 1.00	14.534 0.039	13.909 0.046	13.569 0.047
Annizam 2557215		05 34 36.20	–01 47 13.9	13.969 0.04	12.443 0.022	11.765 0.031	11.532 0.021
Annizam 2166166		05 36 47.81	–01 47 08.8	15.140 1.00	13.175 0.024	12.502 0.032	12.182 0.021
Annizam 2076174		05 36 27.59	–01 46 30.6	16.025 1.00	14.081 0.028	13.422 0.033	13.180 0.029
Annizam 2357151		05 37 29.41	–01 46 25.2	17.279 1.00	15.141 0.046	14.429 0.052	14.245 0.071
Annizam 2485146		05 37 46.01	–01 46 22.2	16.466 1.00	14.648 0.047	13.989 0.079	13.791 0.067
Annizam 2062173		05 36 29.26	–01 46 14.4	16.362 1.00	14.527 0.028	13.872 0.040	13.704 0.032
Annizam 2473146	*	05 37 46.08	–01 46 06.3	16.782 1.00	14.597 0.030	14.088 0.024	13.828 0.036
Annizam 2110161		05 36 58.51	–01 45 22.9	15.516 1.00	13.490 0.027	12.935 0.036	12.570 0.023
Annizam 2286211		05 34 54.43	–01 44 47.3	16.782 0.10	13.381 0.024	12.511 0.032	11.901 0.021
Annizam 1949179		05 36 15.37	–01 44 35.9	16.858 1.00	14.784 0.037	14.265 0.046	14.037 0.042
Annizam 2499221		05 34 24.22	–01 43 42.1	15.965 0.07	14.084 0.026	13.468 0.033	13.192 0.032
Annizam 1926168		05 36 38.75	–01 43 33.6	16.052 1.00	14.072 0.030	13.505 0.038	13.179 0.030
Annizam 2464138	*	05 38 02.89	–01 42 36.9	15.104 1.00	13.124 0.027	12.582 0.024	12.234 0.028
Annizam 2360221		05 34 30.47	–01 41 59.3	16.760 0.10	14.792 0.035	14.060 0.043	13.849 0.043
Annizam 2122146		05 37 31.84	–01 41 26.7	16.513 1.00	14.495 0.032	13.885 0.027	13.577 0.038
Annizam 2684230		05 33 56.47	–01 41 06.1	15.200 0.05	13.474 0.024	12.811 0.035	12.540 0.023
Annizam 2225141		05 37 45.57	–01 41 03.8	15.792 1.00	14.077 0.026	13.429 0.023	13.118 0.029
Annizam 1768200		05 35 32.93	–01 39 50.9	16.486 0.07	14.744 0.032	14.049 0.042	13.828 0.040
Annizam 1719165		05 36 43.34	–01 39 43.5	16.311 1.00	14.342 0.031	13.700 0.038	13.467 0.026
Annizam 1974146		05 37 26.44	–01 39 22.8	16.396 1.00	14.484 0.029	13.935 0.049	13.622 0.052
Annizam 1661163		05 36 45.35	–01 38 35.0	15.911 1.00	14.096 0.027	13.491 0.033	13.263 0.028
Annizam 1840146	*	05 37 21.24	–01 37 34.5	17.623 1.00	15.359 0.058	14.566 0.064	14.187 0.076
Annizam 2556126		05 38 30.63	–01 37 10.9	16.375 0.08	14.959 0.032	14.153 0.033	13.682 0.042
Annizam 1530169		05 36 32.80	–01 37 07.2	14.832 1.00	12.943 0.024	12.293 0.033	12.026 0.023
Annizam 1498173		05 36 24.96	–01 36 53.9	15.796 1.00	14.094 0.028	13.477 0.033	13.226 0.026
Annizam 2456234		05 34 00.38	–01 36 12.1	17.818 0.17	15.414 0.047	14.774 0.067	14.415 0.066
Annizam 2648123		05 38 41.37	–01 35 58.5	14.932 0.04	13.052 0.027	12.424 0.026	12.193 0.029
Annizam 2135131		05 37 59.72	–01 35 36.7	15.157 1.00	13.419 0.028	12.790 0.024	12.538 0.033
Annizam 2254128		05 38 10.73	–01 35 24.5	16.268 1.00	14.314 0.030	13.740 0.033	13.359 0.028
Annizam 1686145		05 37 17.55	–01 35 05.0	16.904 1.00	14.489 0.030	14.014 0.040	13.673 0.048
Annizam 2392237		05 33 59.05	–01 33 50.8	17.263 0.13	14.697 0.039	14.045 0.036	13.627 0.030
Annizam 1685141		05 37 24.05	–01 33 50.3	17.639 1.00	15.274 0.041	14.644 0.073	14.282 0.079
Annizam 2576240		05 33 44.30	–01 33 42.5	14.593 0.04	12.922 0.024	12.261 0.032	11.974 0.019
Annizam 2267124		05 38 17.69	–01 33 25.4	16.533 0.09	14.686 0.031	14.131 0.033	13.674 0.048
Annizam 2331237		05 34 02.39	–01 33 16.0	14.970 0.05	13.333 0.027	12.669 0.032	12.361 0.024
Annizam 2262238		05 34 04.32	–01 31 52.5	16.694 0.10	14.823 0.035	14.139 0.048	13.874 0.043
Annizam 2630244		05 33 35.57	–01 31 32.0	17.140 0.12	14.926 0.033	14.372 0.029	14.083 0.065
Annizam 1843129		05 37 48.15	–01 31 30.6	16.367 1.00	14.535 0.038	13.781 0.029	13.478 0.049
Annizam 1524221		05 35 06.01	–01 31 15.9	15.719 0.05	14.032 0.030	13.448 0.038	13.089 0.030
Annizam 1891125		05 37 55.55	–01 30 23.0	15.700 1.00	14.031 0.028	13.283 0.026	13.052 0.033
Annizam 1897236		05 34 28.37	–01 29 57.8	16.304 0.08	14.478 0.029	13.849 0.037	13.581 0.035

Table A.16. Young star and brown dwarf candidates in the Alnilam field and in DENIS/2MASS (cont.).

Identification		α (J2000)	δ (J2000)	$i \pm \delta i$ (mag)	$J \pm \delta J$ (mag)	$H \pm \delta H$ (mag)	$K_s \pm \delta K_s$ (mag)
Annizam 1460135		05 37 21.98	–01 29 14.0	16.622 1.00	14.209 0.029	13.703 0.035	13.333 0.046
Annizam 1119210		05 35 35.31	–01 28 14.7	16.058 0.06	14.249 0.026	13.553 0.030	13.326 0.031
Annizam 1512232		05 34 53.01	–01 27 31.9	16.490 0.09	14.656 0.032	14.053 0.040	13.645 0.030
Annizam 1142216		05 35 27.87	–01 27 28.8	15.132 0.04	13.477 0.026	12.716 0.031	12.397 0.023
Annizam 1407131		05 37 23.94	–01 27 23.8	14.311 1.00	12.636 0.029	11.952 0.022	11.739 0.024
Annizam 2059245		05 34 08.84	–01 26 53.1	14.883 0.05	13.292 0.026	12.619 0.032	12.361 0.026
Annizam 952149		05 36 45.19	–01 25 45.5	16.710 1.00	14.756 0.036	14.185 0.051	13.794 0.058
Annizam 2168249		05 33 58.22	–01 25 19.9	16.376 0.09	14.568 0.032	13.941 0.037	13.664 0.036
Annizam 875152		05 36 40.12	–01 25 00.6	16.460 1.00	14.558 0.030	13.997 0.036	13.686 0.037
Annizam 1868114		05 38 06.36	–01 24 54.2	17.411 1.00	14.996 0.031	14.417 0.033	14.080 0.060
Annizam 1483120		05 37 38.53	–01 24 27.0	16.632 1.00	14.592 0.031	14.037 0.040	13.625 0.041
Annizam 807149		05 36 40.51	–01 23 38.8	15.037 1.00	12.806 0.022	12.140 0.030	11.873 0.023
Annizam 2574254		05 33 27.39	–01 23 35.0	16.005 0.07	14.040 0.028	13.481 0.026	13.142 0.034
Annizam 699167		05 36 23.21	–01 23 28.0	15.964 1.00	14.047 0.027	13.440 0.033	13.183 0.029
Annizam 1106127	*	05 37 11.43	–01 23 17.8	14.558 1.00	12.983 0.042	12.323 0.051	12.043 0.039
Annizam 2622104		05 39 02.41	–01 22 43.7	15.957 0.06	14.114 0.032	13.384 0.029	12.985 0.023
Annizam 842223		05 35 34.81	–01 22 26.7	16.389 0.07	14.621 0.033	13.937 0.044	13.671 0.040
Annizam 1464112		05 37 43.13	–01 21 22.7	16.129 1.00	14.205 0.028	13.517 0.026	13.220 0.042
Annizam 2636101		05 39 05.13	–01 20 48.0	14.792 0.21	13.227 0.030	12.416 0.027	12.170 0.029
Annizam 1696253		05 34 24.88	–01 20 35.1	17.305 0.13	15.185 0.036	14.423 0.043	14.105 0.043
Annizam 863125		05 36 59.83	–01 20 24.5	14.749 1.00	12.963 0.030	12.075 0.023	11.343 0.019
Annizam 2561101		05 39 00.67	–01 19 56.7	13.656 0.03	12.587 0.056	11.578 0.030	11.338 0.029
Annizam 966118		05 37 09.58	–01 19 44.0	13.942 1.00	12.425 0.024	11.702 0.023	11.508 0.023
Annizam 1143247		05 35 02.85	–01 19 40.1	16.101 0.06	14.417 0.032	13.743 0.035	13.462 0.035
Annizam 1716105		05 38 03.24	–01 19 36.0	15.796 1.00	14.077 0.031	13.401 0.026	13.178 0.035
Annizam 662133		05 36 45.35	–01 19 35.0	15.283 1.00	13.620 0.031	12.915 0.035	12.461 0.024
Annizam 991116		05 37 11.96	–01 19 27.9	16.624 1.00	14.499 0.037	13.868 0.045	13.672 0.048
Annizam 973117		05 37 10.87	–01 19 21.6	14.004 1.00	12.267 0.024	11.619 0.022	11.349 0.023
Annizam 1761256		05 34 18.92	–01 19 16.2	15.920 0.07	14.183 0.033	13.556 0.042	13.196 0.031
Annizam 1711102		05 38 04.33	–01 18 08.7	16.706 1.00	14.538 0.035	13.933 0.026	13.590 0.051
Annizam 1083251		05 35 04.58	–01 18 01.8	15.883 0.05	14.316 0.030	13.591 0.032	13.314 0.030
Annizam 2480098		05 38 56.53	–01 17 58.7	17.472 0.24	15.385 0.050	14.760 0.041	14.424 0.070
Annizam 1113106		05 37 24.34	–01 17 05.6	13.721 1.00	12.154 0.024	11.464 0.023	11.253 0.024
Annizam 1415101	*	05 37 45.42	–01 16 36.3	16.845 1.00	14.795 0.038	14.196 0.037	13.880 0.052
Annizam 2525264		05 33 25.36	–01 16 26.2	16.374 0.09	14.603 0.032	14.026 0.035	13.706 0.045
Annizam 1892098		05 38 17.82	–01 16 23.8	16.185 0.07	14.387 0.032	13.757 0.032	13.458 0.044
Annizam 2118264		05 33 52.47	–01 16 02.7	14.447 0.04	12.849 0.023	12.135 0.030	11.860 0.023
Annizam 1651098		05 38 01.92	–01 15 45.5	14.816 1.00	13.121 0.031	12.547 0.030	12.266 0.037
Annizam 2042095	*	05 38 28.55	–01 14 52.2	16.417 0.08	14.542 0.031	13.886 0.033	13.379 0.030
Annizam 2668267		05 33 15.16	–01 14 22.1	16.831 0.11	14.939 0.035	14.310 0.037	14.011 0.061
Annizam 937262		05 35 10.95	–01 14 14.9	15.160 0.04	13.241 0.024	12.640 0.031	12.358 0.023
Annizam 726099		05 37 00.59	–01 14 01.4	17.262 1.00	14.965 0.044	14.268 0.039	13.733 0.048
Annizam 2611268	*	05 33 18.83	–01 13 55.6	16.387 0.09	14.477 0.033	13.888 0.040	13.559 0.037
Annizam 123132	*	05 36 18.89	–01 13 29.1	16.613 1.00	14.633 0.030	13.993 0.033	13.671 0.036
Annizam 2441092		05 38 55.53	–01 13 22.5	17.393 0.24	15.352 0.053	14.586 0.043	14.299 0.063
Annizam 739094		05 37 01.99	–01 12 52.2	16.076 1.00	14.341 0.036	13.697 0.025	13.406 0.047
Annizam 2404269		05 33 32.51	–01 12 51.9	16.304 0.08	14.013 0.024	13.347 0.027	12.921 0.027
Annizam 1094091		05 37 25.75	–01 12 31.5	15.854 1.00	14.028 0.032	13.341 0.029	13.128 0.033
Annizam 1187092		05 34 53.70	–01 11 24.7	14.581 0.04	12.972 0.022	12.333 0.032	12.046 0.026
Annizam 1535268		05 37 55.09	–01 11 14.2	16.198 1.00	14.248 0.036	13.504 0.029	13.256 0.033
Annizam 727264		05 37 01.05	–01 10 56.6	16.006 1.00	14.043 0.038	13.472 0.036	13.159 0.036
Annizam 1751268	*	05 38 09.46	–01 10 51.2	12.138 1.00	10.592 0.026	9.815 0.024	9.327 0.024
Annizam 1356267		05 37 43.12	–01 10 50.7	16.764 1.00	14.377 0.030	13.776 0.026	13.441 0.039

Table A.16. Young star and brown dwarf candidates in the Alnilam field and in DENIS/2MASS (cont.).

Identification		α (J2000)	δ (J2000)	$i \pm \delta i$ (mag)	$J \pm \delta J$ (mag)	$H \pm \delta H$ (mag)	$K_s \pm \delta K_s$ (mag)
Annizam 860265		05 37 09.92	–01 10 50.4	15.050 1.00	13.179 0.031	12.560 0.027	12.206 0.023
Annizam 1748267	*	05 38 09.24	–01 10 43.2	14.431 1.00	12.635 0.040	11.926 0.043	11.719 0.031
Annizam 1352266		05 37 42.77	–01 10 42.6	16.695 1.00	14.817 0.036	14.101 0.037	13.783 0.050
Annizam 1516266		05 37 53.68	–01 10 30.1	16.078 1.00	14.196 0.027	13.533 0.029	13.276 0.033
Annizam 2463093		05 33 28.76	–01 10 05.6	15.022 0.05	13.867 0.022	13.074 0.024	12.379 0.023
Annizam 1067097		05 35 02.20	–01 09 59.5	16.405 0.07	14.585 0.034	13.923 0.038	13.617 0.039
Annizam 2542267		05 39 02.04	–01 09 49.0	15.820 0.06	14.072 0.027	13.413 0.023	13.243 0.038
Annizam 987099		05 35 07.84	–01 09 28.9	14.478 0.03	12.802 0.021	12.114 0.026	11.912 0.023
Annizam 1634263		05 38 01.01	–01 08 58.4	15.892 1.00	14.006 0.030	13.469 0.030	13.136 0.034
Annizam 1183261		05 37 30.63	–01 08 53.3	15.220 1.00	13.479 0.030	12.805 0.026	12.578 0.030
Annizam 1024259		05 37 19.79	–01 08 46.3	13.757 1.00	12.175 0.027	11.459 0.027	11.252 0.024
Annizam 2663266		05 39 09.88	–01 08 45.9	14.595 0.20	12.871 0.027	12.163 0.024	11.962 0.024
Annizam 1724262		05 38 06.74	–01 08 17.8	16.640 1.00	14.219 0.028	13.638 0.027	13.284 0.033
Annizam 1601261		05 37 58.19	–01 07 51.3	17.945 1.00	15.459 0.054	14.688 0.048	14.479 0.083
Annizam 1714099		05 34 19.87	–01 07 44.1	15.955 0.07	14.204 0.032	13.551 0.036	13.302 0.030
Annizam 678246		05 36 54.26	–01 07 36.3	16.502 1.00	14.520 0.032	13.857 0.036	13.671 0.036
V993 Ori	*	05 37 59.04	–01 05 52.8	12.183 1.00	10.846 0.027	9.955 0.024	9.355 0.027
Annizam 806242		05 37 00.28	–01 05 48.6	16.605 1.00	14.568 0.041	14.109 0.052	13.694 0.058
Annizam 455211		05 36 28.32	–01 05 35.6	16.107 1.00	14.324 0.028	13.674 0.030	13.349 0.033
Annizam 638230		05 36 45.56	–01 05 20.0	17.099 1.00	15.017 0.039	14.341 0.052	13.797 0.051
Annizam 2000102		05 34 02.33	–01 05 12.9	16.977 0.12	15.137 0.040	14.479 0.061	14.117 0.049
Annizam 1080242		05 37 16.28	–01 03 36.4	16.816 1.00	14.634 0.032	14.058 0.045	13.573 0.042
[SE2005] 105		05 35 25.62	–01 02 55.8	14.138 0.03	12.450 0.024	11.839 0.033	11.526 0.019
Annizam 1089239		05 37 15.33	–01 02 53.3	17.905 1.00	15.120 0.049	14.629 0.064	14.275 0.076
[SE2005] 13		05 33 53.35	–01 02 12.9	16.191 0.08	14.409 0.036	13.747 0.048	13.417 0.037
Annizam 1114236		05 37 14.66	–01 01 50.5	14.536 1.00	12.602 0.028	12.029 0.027	11.708 0.023
Annizam 1238122		05 35 02.58	–01 01 16.3	15.984 0.06	14.378 0.027	13.589 0.036	13.408 0.036
Annizam 863141		05 35 36.78	–01 00 53.9	15.931 0.05	14.007 0.035	13.537 0.047	13.138 0.033
Annizam 1072228		05 37 06.26	–01 00 14.9	14.706 1.00	13.066 0.028	12.389 0.026	12.061 0.027
Annizam 2000249		05 38 17.30	–01 00 10.5	14.742 0.04	12.937 0.026	12.297 0.023	11.930 0.027
Annizam 2540107		05 33 31.07	–00 59 35.2	17.470 0.14	15.440 0.055	14.725 0.055	14.452 0.081
Annizam 798196	*	05 36 27.54	–00 59 20.5	14.960 1.00	13.314 0.023	12.628 0.034	12.408 0.022
Annizam 817183		05 36 15.92	–00 58 30.8	15.887 1.00	14.041 0.021	13.475 0.046	13.154 0.022
Annizam 1487236		05 37 35.33	–00 58 22.1	17.710 1.00	15.016 0.033	14.520 0.044	14.134 0.057
Annizam 1703119		05 34 33.53	–00 58 19.5	12.273 0.03	10.762 0.022	9.814 0.032	9.536 0.021
Annizam 1380230		05 37 23.50	–00 57 24.0	16.664 1.00	14.661 0.039	14.069 0.052	13.769 0.045
Annizam 1111147		05 35 32.23	–00 56 37.4	16.437 0.07	14.662 0.032	14.021 0.045	13.766 0.039
Annizam 1010201		05 36 36.42	–00 56 20.9	17.312 1.00	15.286 0.039	14.678 0.057	14.194 0.054
Annizam 1628234		05 37 40.69	–00 56 11.5	15.938 1.00	14.306 0.030	13.725 0.026	13.381 0.037
Annizam 1449229		05 37 25.25	–00 56 08.3	16.929 1.00	14.445 0.034	13.820 0.023	13.463 0.037
Annizam 1806123		05 34 31.69	–00 55 46.7	16.216 0.08	14.377 0.027	13.764 0.037	13.542 0.032
Annizam 1352223		05 37 14.64	–00 55 42.6	16.873 1.00	14.926 0.034	14.375 0.060	13.964 0.061
Annizam 2564113		05 33 35.14	–00 55 35.1	15.086 0.05	13.693 0.026	12.718 0.024	12.159 0.024
Annizam 1207210		05 36 53.16	–00 54 42.6	15.997 1.00	14.383 0.028	13.753 0.033	13.456 0.033
Annizam 2406116		05 33 48.84	–00 54 26.0	17.029 0.12	14.670 0.033	14.113 0.047	13.679 0.036
Annizam 1911124		05 34 27.01	–00 54 22.7	15.730 0.07	14.129 0.027	13.478 0.042	13.115 0.030
Annizam 2229241		05 38 22.89	–00 54 08.2	17.861 1.00	15.423 0.046	14.793 0.049	14.539 0.090
Annizam 1289211		05 36 57.26	–00 53 44.1	16.324 1.00	14.633 0.035	13.903 0.041	13.655 0.033
Annizam 1173197		05 36 36.26	–00 53 27.9	16.485 1.00	14.511 0.031	13.921 0.037	13.637 0.035
Annizam 2112123		05 34 14.61	–00 52 59.3	16.149 0.08	14.179 0.027	13.582 0.037	13.233 0.029
Annizam 1763132		05 34 45.70	–00 52 23.6	16.424 0.09	14.519 0.043	13.976 0.050	13.622 0.052
Annizam 2270238		05 38 20.90	–00 51 58.0	15.267 0.05	13.464 0.026	12.899 0.029	12.599 0.030
Annizam 1982232		05 37 57.46	–00 51 56.0	15.962 1.00	14.200 0.032	13.543 0.029	13.272 0.033

Table A.16. Young star and brown dwarf candidates in the Alnilam field and in DENIS/2MASS (cont.).

Identification		α (J2000)	δ (J2000)	$i \pm \delta i$ (mag)	$J \pm \delta J$ (mag)	$H \pm \delta H$ (mag)	$K_s \pm \delta K_s$ (mag)
Annizam 2518241		05 38 39.44	–00 51 40.8	16.848 0.10	14.416 0.028	13.774 0.033	13.342 0.033
[SE2005] 120	*	05 35 38.51	–00 51 11.5	15.376 0.04	13.338 0.024	12.569 0.029	12.301 0.018
Annizam 2649241		05 38 47.24	–00 50 42.0	14.114 0.03	12.195 0.026	11.606 0.026	11.280 0.019
Annizam 1643214		05 37 14.78	–00 49 31.8	14.009 1.00	12.129 0.024	11.386 0.023	11.038 0.021
Annizam 1444191		05 36 31.14	–00 48 29.7	17.098 1.00	15.293 0.055	14.347 0.049	13.714 0.036
Annizam 2419126		05 34 02.41	–00 48 23.1	17.369 0.14	15.364 0.049	14.778 0.061	14.280 0.064
Annizam 1569198		05 36 44.76	–00 47 12.6	16.501 1.00	14.345 0.024	13.709 0.040	13.433 0.024
[SE2005] 126	*	05 35 43.63	–00 46 46.5	17.940 0.15	15.768 0.066	15.042 0.082	14.608 0.068
Annizam 1724208		05 37 06.26	–00 46 40.6	16.293 1.00	14.498 0.038	13.918 0.031	13.629 0.052
Annizam 1619161		05 35 38.02	–00 46 34.2	16.634 0.08	14.595 0.044	14.012 0.047	13.706 0.044
Annizam 1805154		05 35 19.55	–00 45 08.4	15.933 0.05	14.058 0.026	13.459 0.036	13.202 0.030
Annizam 1680196		05 36 42.89	–00 45 08.2	15.067 1.00	13.198 0.024	12.351 0.028	12.085 0.019
Annizam 1692195		05 36 41.75	–00 44 52.0	17.275 1.00	14.890 0.036	14.219 0.045	13.982 0.039
Annizam 2650232		05 38 31.44	–00 44 44.0	17.060 0.12	15.034 0.034	14.321 0.035	14.051 0.059
[SE2005] 67		05 34 44.91	–00 43 50.2	15.560 0.06	13.131 0.024	12.334 0.026	11.994 0.021
Annizam 2388137		05 34 23.63	–00 43 09.0	16.424 0.09	14.541 0.028	13.925 0.033	13.586 0.036
Annizam 1909156		05 35 21.16	–00 43 01.7	17.030 0.09	14.693 0.031	14.136 0.042	13.687 0.037
Annizam 2504135		05 34 14.23	–00 42 43.8	16.255 0.08	14.336 0.031	13.701 0.035	13.452 0.030
Annizam 1823193		05 36 39.81	–00 42 29.7	15.148 1.00	13.216 0.024	12.565 0.030	12.298 0.019
Annizam 2101211		05 37 25.90	–00 42 14.3	16.138 1.00	14.112 0.032	13.517 0.041	13.219 0.036
Annizam 2135212		05 37 29.09	–00 42 04.0	16.664 1.00	14.578 0.024	13.981 0.036	13.687 0.044
[SE2005] 99		05 35 22.44	–00 40 03.2	14.775 0.04	13.155 0.024	12.360 0.030	11.970 0.023
Annizam 2140198		05 36 57.07	–00 38 12.1	16.797 1.00	14.335 0.027	13.652 0.036	13.394 0.030
Annizam 2282155		05 35 07.91	–00 37 43.3	16.277 0.06	14.316 0.024	13.715 0.037	13.391 0.029
Annizam 2690220		05 38 07.37	–00 37 37.3	16.935 1.00	15.185 0.050	14.499 0.046	14.071 0.061
Annizam 2291204		05 37 15.97	–00 37 21.0	16.318 1.00	14.495 0.039	13.938 0.025	13.576 0.045
[SE2005] 71	*	05 34 48.00	–00 37 17.9	14.901 0.05	13.160 0.022	12.517 0.025	12.308 0.023
Annizam 2194198		05 36 57.32	–00 37 17.4	16.438 1.00	14.557 0.032	13.839 0.039	13.618 0.029
Annizam 2477210		05 37 34.28	–00 36 12.1	15.096 1.00	13.426 0.028	12.828 0.027	12.586 0.027
Annizam 2357160		05 35 19.63	–00 35 09.0	17.251 0.11	14.988 0.040	14.437 0.049	14.013 0.045
Annizam 2619195		05 36 58.52	–00 29 59.4	14.188 1.00	12.204 0.024	11.541 0.030	11.298 0.021

Table A.17. Young star and brown dwarf candidates in the Mintaka field and in DENIS/2MASS.

Identification		α (J2000)	δ (J2000)	$i \pm \delta i$ (mag)	$J \pm \delta J$ (mag)	$H \pm \delta H$ (mag)	$K_s \pm \delta K_s$ (mag)
Mantaqah 2389165		05 32 42.57	–00 56 20.5	17.036 0.11	14.988 0.035	14.474 0.052	14.011 0.058
Mantaqah 2668148		05 33 35.14	–00 55 35.1	15.176 0.05	13.693 0.026	12.718 0.024	12.159 0.024
Mantaqah 2473149		05 33 24.52	–00 53 23.9	14.933 0.05	13.358 0.026	12.737 0.027	12.430 0.024
Mantaqah 1995178		05 32 05.67	–00 51 09.7	15.705 0.05	14.021 0.022	13.357 0.023	13.139 0.037
Mantaqah 2691223		05 29 58.45	–00 50 50.3	18.219 1.00	15.521 0.056	14.783 0.056	14.503 0.086
Mantaqah 1970186		05 31 45.96	–00 50 35.0	14.409 0.03	12.785 0.024	12.103 0.027	11.836 0.026
Mantaqah 2445218		05 30 19.69	–00 49 59.7	16.398 1.00	14.254 0.032	13.532 0.032	13.225 0.034
Mantaqah 2041159	*	05 32 48.68	–00 49 44.8	14.644 0.04	12.969 0.033	12.396 0.040	12.088 0.033
Mantaqah 1876186		05 31 47.79	–00 49 02.9	15.055 0.04	13.242 0.026	12.395 0.024	12.139 0.019
Mantaqah 1957161		05 32 43.46	–00 48 44.5	15.097 0.05	13.341 0.026	12.776 0.029	12.486 0.032
Mantaqah 1832184		05 31 52.38	–00 48 25.2	13.479 0.03	11.539 0.023	10.653 0.024	10.317 0.021
[SE2005] 22		05 34 02.41	–00 48 23.1	17.369 0.14	15.364 0.049	14.778 0.061	14.280 0.064
Mantaqah 1778177		05 32 06.21	–00 47 32.7	16.783 0.09	14.658 0.037	14.060 0.037	13.779 0.047
Mantaqah 2541226		05 29 58.50	–00 47 21.0	15.409 1.00	13.448 0.027	12.760 0.025	12.524 0.027
Mantaqah 1730181		05 31 59.23	–00 46 46.3	17.230 0.11	15.118 0.030	14.454 0.046	14.115 0.065
Mantaqah 2423225		05 30 05.61	–00 46 22.0	17.231 1.00	14.576 0.035	14.067 0.043	13.683 0.052
[SE2005] 44	*	05 34 23.63	–00 43 09.0	16.424 0.09	14.541 0.028	13.925 0.033	13.586 0.036
[SE2005] 32		05 34 14.23	–00 42 43.8	16.255 0.08	14.336 0.031	13.701 0.035	13.452 0.030
Mantaqah 2216132	*	05 33 50.20	–00 42 39.7	15.160 0.05	13.253 0.027	12.639 0.036	12.352 0.024
Mantaqah 1493159		05 32 36.66	–00 41 07.2	17.026 0.11	14.789 0.033	14.194 0.042	13.817 0.051
Mantaqah 2426235		05 29 47.62	–00 41 02.3	16.087 1.00	14.287 0.030	13.549 0.028	13.327 0.039
Mantaqah 2265235		05 29 57.46	–00 39 51.8	14.575 1.00	12.962 0.030	12.322 0.035	12.081 0.033
Mantaqah 2232234		05 29 59.49	–00 39 38.6	16.994 1.00	15.222 0.040	14.499 0.032	14.163 0.062
Mantaqah 2533239		05 29 35.51	–00 39 38.4	15.901 0.05	14.183 0.029	13.464 0.028	13.215 0.039
Mantaqah 2263235		05 29 56.96	–00 39 38.0	16.875 1.00	14.829 0.040	14.248 0.040	13.896 0.046
Mantaqah 1592144		05 33 03.48	–00 39 16.8	16.885 0.10	14.917 0.036	14.291 0.039	13.959 0.060
Mantaqah 1287190		05 31 45.86	–00 39 05.1	16.193 0.07	14.200 0.029	13.463 0.029	13.111 0.036
Mantaqah 1284170		05 32 15.60	–00 39 00.1	16.218 0.07	14.263 0.028	13.199 0.027	12.519 0.024
Mantaqah 1992129		05 33 43.23	–00 38 57.9	14.724 0.05	12.909 0.024	12.287 0.030	12.015 0.019
Mantaqah 2037232		05 30 12.92	–00 38 41.4	15.099 1.00	13.403 0.026	12.804 0.022	12.555 0.031
Mantaqah 1218172		05 32 11.42	–00 38 03.6	14.518 0.03	12.941 0.026	12.322 0.027	11.996 0.023
Mantaqah 1592138		05 33 11.31	–00 37 41.0	15.099 0.05	13.148 0.029	12.495 0.023	12.147 0.024
Mantaqah 2078236		05 30 06.09	–00 37 30.4	16.961 1.00	14.972 0.035	14.351 0.046	14.011 0.053
Mantaqah 1543139		05 33 07.23	–00 37 29.5	14.880 0.05	13.298 0.023	12.612 0.023	12.368 0.024
Mantaqah 1131192		05 31 44.10	–00 36 20.9	15.782 0.05	14.043 0.030	13.370 0.030	13.149 0.034
Mantaqah 1080171		05 32 11.58	–00 35 43.7	17.412 0.12	15.212 0.043	14.478 0.036	14.115 0.075
Mantaqah 2147240		05 29 56.13	–00 35 41.6	17.497 1.00	15.142 0.042	14.433 0.040	14.000 0.070
Mantaqah 1076190		05 31 47.88	–00 35 36.6	16.730 0.09	14.735 0.036	14.202 0.037	13.842 0.060
Mantaqah 2473115		05 34 30.44	–00 35 03.1	16.255 0.08	14.536 0.034	13.853 0.041	13.595 0.035
Mantaqah 1368135		05 33 05.17	–00 33 59.8	14.568 0.04	12.962 0.035	12.329 0.039	12.023 0.035
Mantaqah 2385113	*	05 34 26.30	–00 33 44.4	14.574 0.04	12.675 0.052	11.998 0.060	11.738 0.054
Mantaqah 2349114		05 34 23.81	–00 33 40.7	12.375 0.03	10.760 0.021	9.829 0.030	9.546 0.019
Mantaqah 849188		05 31 52.73	–00 31 57.8	16.671 0.08	14.771 0.033	14.069 0.035	13.791 0.051
Mantaqah 865162		05 32 17.88	–00 31 41.5	14.981 0.04	13.124 0.028	12.179 0.027	11.677 0.023
Mantaqah 2507108		05 34 39.13	–00 31 03.4	14.362 0.04	12.554 0.024	11.968 0.029	11.689 0.024
Mantaqah 1441116		05 33 26.74	–00 28 28.7	16.171 0.08	14.233 0.023	13.663 0.036	13.329 0.030
Mantaqah 1761250		05 30 10.17	–00 28 01.5	17.096 1.00	14.960 0.040	14.370 0.042	14.078 0.065
Mantaqah 1787109		05 33 52.77	–00 27 51.3	15.785 0.07	14.033 0.026	13.438 0.031	13.120 0.032
Mantaqah 1547248		05 30 24.59	–00 27 29.8	14.650 1.00	13.029 0.024	12.368 0.024	12.143 0.026
Mantaqah 746137		05 32 34.34	–00 27 01.8	14.757 0.05	13.050 0.024	12.399 0.025	12.209 0.026
Mantaqah 761135		05 32 36.02	–00 26 59.2	16.240 0.08	14.164 0.028	13.655 0.027	13.148 0.033
Mantaqah 568156		05 32 15.50	–00 26 37.5	15.152 0.04	13.421 0.028	12.818 0.031	12.559 0.031

Table A.17. Young star and brown dwarf candidates in the Mintaka field and in DENIS/2MASS (cont.).

Identification		α (J2000)	δ (J2000)	$i \pm \delta i$ (mag)	$J \pm \delta J$ (mag)	$H \pm \delta H$ (mag)	$K_s \pm \delta K_s$ (mag)
Mantaqah 570147		05 32 21.37	–00 25 52.6	15.955 0.06	14.106 0.040	13.492 0.047	13.121 0.041
Mantaqah 557212		05 31 40.70	–00 25 48.8	17.160 0.11	14.926 0.039	14.066 0.026	13.347 0.043
Mantaqah 1787105		05 33 55.73	–00 25 24.7	15.264 0.05	13.638 0.027	12.771 0.040	12.164 0.026
Mantaqah 1361109		05 33 26.24	–00 25 18.9	15.441 0.06	13.483 0.023	12.903 0.025	12.641 0.027
Mantaqah 2099258		05 29 43.30	–00 24 55.2	16.098 0.05	14.394 0.029	13.806 0.032	13.490 0.041
Mantaqah 1606104		05 33 44.10	–00 24 37.7	13.778 0.03	11.967 0.024	11.313 0.031	11.020 0.021
Mantaqah 2074101		05 34 16.21	–00 24 27.3	14.253 0.04	12.452 0.021	11.819 0.031	11.564 0.022
Mantaqah 2656098		05 34 55.55	–00 24 24.5	16.441 0.09	14.476 0.028	13.825 0.035	13.565 0.030
Mantaqah 352154		05 32 10.65	–00 23 13.2	16.118 0.06	14.158 0.045	13.560 0.048	13.300 0.047
Mantaqah 821105		05 32 53.30	–00 21 29.2	15.869 0.07	14.004 0.029	13.458 0.022	13.144 0.027
Mantaqah 941101	*	05 33 01.97	–00 20 59.3	13.396 0.03	11.989 0.024	11.115 0.025	10.551 0.024
Mantaqah 325118		05 32 19.49	–00 20 30.1	16.662 0.09	14.669 0.033	14.158 0.038	13.788 0.055
Mantaqah 1926266	*	05 29 52.34	–00 20 23.2	16.444 1.00	14.344 0.029	13.780 0.034	13.475 0.039
Mantaqah 161138	*	05 32 07.55	–00 19 56.3	15.241 0.04	13.350 0.026	12.770 0.027	12.476 0.023
Mantaqah 1408095		05 33 33.93	–00 19 50.9	16.181 0.08	14.414 0.033	13.750 0.030	13.530 0.048
Mantaqah 104142	*	05 32 04.69	–00 19 18.9	17.955 0.17	15.156 0.042	14.338 0.045	14.194 0.077
Mantaqah 67144	*	05 32 03.03	–00 18 51.4	15.159 0.04	12.942 1.000	12.866 0.026	12.624 0.027
Mantaqah 125248		05 31 52.67	–00 18 43.5	16.599 0.08	14.773 0.032	14.145 0.037	13.705 0.049
Mantaqah 115248		05 31 53.31	–00 18 40.1	17.243 0.11	14.775 0.033	14.202 0.043	13.804 0.054
Mantaqah 2270090		05 34 31.71	–00 18 00.0	16.096 0.08	14.209 0.026	13.597 0.038	13.321 0.027
Mantaqah 2299090		05 34 33.69	–00 17 58.8	15.943 0.07	14.173 0.031	13.571 0.038	13.289 0.035
Mantaqah 378267		05 32 25.57	–00 17 34.3	14.170 0.04	12.614 0.023	11.897 0.022	11.555 0.023
Mantaqah 718263		05 32 47.87	–00 16 23.8	15.741 0.06	14.049 0.024	13.451 0.029	13.173 0.033
Mantaqah 808261		05 32 53.64	–00 15 51.8	17.073 0.11	14.648 0.029	14.038 0.025	13.686 0.039
Mantaqah 1967266		05 34 11.24	–00 15 51.0	17.925 0.17	15.417 0.045	14.861 0.060	14.499 0.069
Mantaqah 331115		05 31 40.47	–00 15 34.6	17.222 0.11	14.965 0.039	14.383 0.050	14.016 0.051
Mantaqah 1982094	*	05 29 48.63	–00 15 33.5	13.426 1.00	11.849 0.024	11.045 0.024	10.779 0.023
Mantaqah 528246		05 32 32.45	–00 14 17.9	15.948 0.07	14.308 0.026	13.687 0.022	13.356 0.037
Mantaqah 1968098		05 29 50.50	–00 13 21.8	14.504 1.00	12.726 0.026	12.162 0.026	11.880 0.025
Mantaqah 467208		05 32 14.90	–00 11 03.0	15.122 0.04	13.238 0.024	12.624 0.022	12.335 0.026
Mantaqah 1429252		05 33 31.18	–00 10 44.4	16.672 0.10	14.479 0.027	13.880 0.030	13.579 0.041
Mantaqah 2615100		05 29 08.64	–00 10 27.7	16.819 0.08	14.843 0.030	14.226 0.046	13.971 0.054
Mantaqah 1084244		05 33 05.56	–00 10 08.7	17.501 0.14	15.149 0.040	14.553 0.048	14.212 0.070
Mantaqah 802230		05 32 41.53	–00 09 23.7	16.340 0.08	14.315 0.028	13.622 0.022	13.364 0.037
Mantaqah 558158		05 31 46.31	–00 09 20.5	16.516 0.08	14.388 0.032	13.835 0.040	13.462 0.041
Mantaqah 767217		05 32 30.97	–00 07 42.5	15.715 0.06	14.049 0.028	13.411 0.035	13.153 0.027
Mantaqah 1883107		05 30 00.64	–00 08 32.5	16.044 1.00	14.053 0.026	13.476 0.026	13.240 0.040
Mantaqah 2522254		05 34 42.01	–00 06 20.7	15.799 0.07	14.147 0.028	13.621 0.035	13.228 0.029
Mantaqah 2326250		05 34 26.29	–00 04 47.9	15.973 0.07	14.175 0.027	13.500 0.037	13.262 0.028
Mantaqah 976215		05 32 37.87	–00 04 39.0	16.138 0.07	14.257 0.032	13.649 0.023	13.323 0.037
Mantaqah 2174236		05 34 00.81	+00 02 13.2	15.122 0.05	13.457 0.030	12.699 0.026	12.170 0.024
Mantaqah 2406121		05 29 43.51	+00 02 58.2	15.396 0.04	14.004 0.027	13.373 0.032	13.149 0.037
Mantaqah 1601217		05 33 04.21	+00 03 27.2	15.524 0.06	13.476 0.026	12.834 0.026	12.433 0.023
Mantaqah 1291180		05 32 00.17	+00 03 34.2	15.040 0.04	13.377 0.024	12.806 0.022	12.513 0.023
Mantaqah 2327130		05 30 00.73	+00 06 44.2	15.811 1.00	14.019 0.036	13.407 0.026	13.137 0.038
Mantaqah 1706194		05 32 28.23	+00 09 37.7	16.097 0.07	14.299 0.051	13.757 0.076	13.414 0.075
Mantaqah 2036210		05 33 08.27	+00 11 26.6	16.290 0.08	14.622 0.042	14.078 0.044	13.648 0.050
Mantaqah 1813174		05 31 47.19	+00 12 05.9	14.860 0.04	13.277 0.029	12.666 0.031	12.347 0.027
Mantaqah 2659135		05 29 55.57	+00 13 31.8	14.444 1.00	12.723 0.040	12.148 0.037	11.838 0.036
Mantaqah 2657142		05 30 10.49	+00 16 46.5	17.124 1.00	14.606 0.039	13.676 0.024	12.901 0.030

Table A.18. Remarkable fore- and background objects in the comparison fields.

Name		α (J2000)	δ (J2000)	V_T (mag)	K_s (mag)	Remarks	References
TYC 5945 228 1		06 09 19.74	–21 04 36.2	11.16±0.07	5.66±0.03	IRAS	*cs08*
Albus 1	*	06 06 13.39	–20 21 07.3	11.80±0.14	12.76±0.03	bright He-B subdwarf	CS07, VKAS07
19 Lep		06 07 41.64	–19 09 57.1	5.480±0.009	1.05±0.06	M1III, var., d = 220±30 pc	Ad26
CSS 205		06 15 33.16	–18 42 16.6	11.63±0.10	5.850±0.017	IRAS, S-type star	St84
HD 43429		06 15 17.72	–18 28 37.8	6.102±0.009	3.6±0.2	K1III, d = 64±3 pc	Ha76
IRAS 06101–1659		06 12 20.36	–16 59 52.4	10.97±0.05	4.885±0.018	IRAS	*cs08*
θ Lep		06 06 09.32	–14 56 06.9	4.669±0.009	4.52±0.02	A0V, d = 52.2±1.9 pc	MR50
TYC 5360 681 1	*	05 58 16.46	–14 43 31.8	10.98±0.06	5.22±0.02	IRAS	*cs08*
η Lep		05 56 24.29	–14 10 03.7	3.738±0.009	2.90±0.01	F1V, d = 15.04±0.17 pc	AJ17
BD–13 1293	*	05 53 38.44	–13 27 50.5	9.68±0.02	3.2±0.3	IRAS	*cs08*
TYC 5361 817 1		06 01 33.94	–13 16 02.8	12.21±0.14	10.33±0.02	μ = 121±2 mas a^{-1}	*cs08*
BD–12 1344		06 00 56.63	–12 04 45.4	10.50±0.04	4.65±0.02	IRAS	*cs08*
TYC 5356 927 1		05 58 07.67	–11 27 25.2	10.68±0.05	9.18±0.02	μ = 155±2 mas a^{-1}	*cs08*

Table A.19. Reference abbreviations used in this Appendix.

Abbreviation	Reference	Abbreviation	Reference
Sc1886	Schoenfeld 1886	Re88	Renson 1988
Ha1904	Hartmann 1904	Ko89	Kogure et al. 1989
Wo1904	Wolf 1904	Wi89	Wiramihardja et al. 1989
Fr09	Frost 1909	DHu91	Dong & Hu 1991
Da16	Daniel 1916	ML91	Morrell & Levato 1991
AJ17	Adams & Joy 1917	Wi91	Wiramihardja et al. 1991
CP18	Cannon & Pickering 1918–1924	Gr92	Grillo et al. 1992
Ad26	Adams et al. 1926	MMR92	Martín, Magazzù & Rebolo 1992
MKK43	Morgan, Keenan & Kellman 1943	Ou92	Oudmaijer et al. 1992
Ne43	Neubauer 1943	PPR92	Pallavicini, Pasquini & Randich 1992
KH50	Keenan & Hynek 1950	GC93	Gray & Corbally 1993
MR50	Morgan & Roman 1950	McD94	McDowell 1994
Sh52	Sharpless 1952	Co95	Codella et al. 1995
HM53	Haro & Moreno 1953	Ne95	Nesterov et al. 1995
MH55	Mannino & Humblet 1955	YC94	Yun & Clemens 1994
Ro55	Roman 1955	Al96	Alcalá et al. 1996
AK59	Artyukhina & Karimova 1959	No97	Nordström et al. 1997
GBT61	Giclas, Burnham & Thomas 1961	PG97	Paunzen & Gray 1997
Bi65	Bidelman 1965	Pe97	Perryman et al. 1997
CB66	Crawford & Barnes 1966	Yun97	Yun et al. 1997
Lee68	Lee 1968	AJ98	Andrillat & Jaschek 1998
Ko69	Kostyakova 1969	Høg98	Høg et al. 1998
Sa71	Sanduleak 1971	Na99	Nakano et al. 1999
SCh71	Schild & Chaffee 1971	WL99	Woolf & Lambert 1999
SCo71	Schild & Cowley 1971	Al00	Alcalá et al. 2000
BC72	Bernacca & Ciatti 1972	BM00	Bhatt & Manoj 2000
Gr72	Griffin 1972	Ja00	Jarret et al. 2000
HR72	Herbig & Rao 1972	Pa01	Paunzen 2001
Ba74	Bartaya 1974	Pa01a	Paunzen et al. 2001
Sh74	Sharpless 1974	Ue01	Ueda et al. 2001
Gu76	Guetter 1976	Ha02	Harvin et al. 2002
Ha76	Hawarden 1976	OGP02	Ogura, Sugitani & Pickles 2002
WH77	Warren & Hesser 1977	Bé03a	Béjar et al. 2003a
AL77	Abt & Levato 1977	Bé03b	Béjar, Caballero & Rebolo 2003b
SS77	Stephenson & Sanduleak 1977	Kr03	Kraemer et al. 2003
CBD'I78	Ciatti, Bernacca & D'Innocenzo 1978	MMB03	Maheswar, Manoj & Bhatt 2003
WH78	Warren & Hesser 1978	SG03	Salim & Gould 2003
Gu79	Guetter 1979	Vi03	Vieira et al. 2003
Bo81	Borra 1981	Ál04	Álvarez et al. 2004
Gu81	Guetter 1981	No04	Nordström et al. 2004
MacC82	MacConnell 1982	Br05	Briceño et al. 2005
Gi83	Gieseking 1983	SE05	Scholz & Eislöffel (2005)
Me83	Mermilliod 1983	TMW05	Templeton, Mattei & Willson 2005
No84	North 1984	He06	Hernández et al. 2006
St84	Stephenson 1984	McG06	McGehee 2006
Eg85	Eggen 1985	Ku06	Kudryavtsev et al. 2006
St86	Stephenson 1986	CS07	Caballero & Solano 2007
Bo87	Bohlender et al. 1987	Ma07	Marilli et al. (2007)
Bo88	Bopp 1988	TMS07	Terrell, Munari & Siviero 2007
BLM88	Bois, Lanning & Mochnaki 1988	VKAS07	Vennes, Kawka & Allyn Smith 2007
DK88	Downes & Keyes 1988	*cs08*	this work

List of Objects

‘Pleiades’ on page 1
‘ρ Ophiuchi’ on page 1
‘Chamaeleon’ on page 1
‘Taurus-Auriga’ on page 1
‘Orion Nebula Cluster’ on page 1
‘σ Orionis’ on page 1
‘Ori OB1b’ on page 1
‘Ori OB1’ on page 2
‘Alnitak’ on page 2
‘Alnilam’ on page 2
‘Mintaka’ on page 2
‘L1630’ on page 2
‘Flame Nebula’ on page 2
‘Horsehead Nebula’ on page 2
‘Collinder 70’ on page 2
‘Berkeley 20’ on page 2
‘NGC 1981’ on page 2
‘NGC 1990’ on page 2
‘Ced 55h’ on page 2
‘VV Ori AB’ on page 2
‘HD 40071’ on page 4
‘HD 40355 AB’ on page 4
‘IC 434’ on page 5
‘V1299 Ori’ on page 5
‘[OS98] 40B’ on page 5
‘Kiso A–0904 62’ on page 6
‘Ruber 1’ on page 7
‘Ruber 2’ on page 7
‘4C–01.06’ on page 8
‘2MASS J05380010–0122377’ on page 8
‘19 Lep’ on page 8
‘HD 43429’ on page 8
‘θ Lep’ on page 8
‘η Lep’ on page 8
‘Albus 1’ on page 8
‘CSS 205’ on page 8
‘[SE2005] 126’ on page 9
‘Mantaqah 2691223’ on page 9